



UNIVERSITI
KEBANGSAAN
MALAYSIA
The National University
of Malaysia

NARATIF BAHARU UKM
UNIVERSITI
ATAN KITA



9TH REGIONAL CONFERENCE ON
MOLECULAR MEDICINE

RCMM

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**20
23**

Precision Medicine in Healthcare

14-15 OCT
2023

🕒 08:00 a.m. - 05:00 p.m. MYT

📍 Bangi Resort Hotel, Bangi, Selangor

**ABSTRACT
BOOK**

ORAL

O1

Elucidation of Epstein-Barr Virus (EBV) Encoded Latent Membrane Protein-1 (LMP-1) 30 bp Deletion Mutation Effect on Immune Checkpoint Regulation in EBV-Associated Nasopharyngeal Carcinoma (NPC) and Non-NPC Cancers

Naveed Ahmed, Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains Malaysia

O2

High Throughput DIA-NN Proteomics Identifies Altered Extracellular Protein Network in CRISPR-Targeted AGR2 of Breast Cancer Cells

Syazalina Zahari, UKM Medical Molecular Biology Institute (UMBI)

O3

Lipidome Analysis and Potential Lipid Biomarkers in Glioblastoma Tissue Using Untargeted Lipidomic Analysis

Khairunnisa Abdul Rashid, Department of Biomedical Imaging, Faculty of Medicine, University of Malaya

O4

TCR Profiling of Cytotoxic T cells Targeting Neoantigens from Colorectal Cancer Patients

Nurul Ainaa' Adilah Rus Bakaruraini, UKM Medical Molecular Biology Institute (UMBI)

O5

Lutein Inhibits the Triple Negative Breast Cancer Cell Progression by Suppressing the Protumour Effects of Tumour-Associated Macrophages

Redzyque Ramza Ramli, Department of Chemical Pathology, School of Medical Sciences, Universiti Sains Malaysia

O6

Super Enhancer Upstream Regulators Drive KLF6-PDGFB Expression and mTORC1 Pathway Activation in Renal Carcinoma

Nurul Nadia Mohamad Zamberi, UKM Medical Molecular Biology Institute (UMBI)

O7

Predictive Modelling of Cardiovascular Risk in Type 2 Diabetes Mellitus Patients: A K Nearest Neighbours Approach

Kee Ooi Ting, UKM Medical Molecular Biology Institute (UMBI)

O8

Alteration of Gut Microbiota in Systemic Lupus Erythematosus (SLE)

Cham Qin Shi, Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia

O9

Gene Expression Profiling of the Extracellular Vesicles From the Insulin-Resistant Adipocytes

Vicneswary Dorairaj, UKM Medical Molecular Biology Institute (UMBI)

ORAL

O10

Functional Impact of Selected DNA Repair Gene Polymorphisms: Insights From In Silico Analyses

Hilary Sito Pei-Ching, UKM Medical Molecular Biology Institute (UMBI)

O11

Characterisation of Extracellular Vesicle Released From Hepatocytes With Excessive Lipid Accumulation

Muhammad Nafiz Haidi Adrus, UKM Medical Molecular Biology Institute (UMBI)

O12

Metabolic Signatures in Obesity: Discovery of Promising Independent Biomarkers

Anis Adibah Osman, UKM Medical Molecular Biology Institute (UMBI)

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Elucidation of Epstein-Barr virus (EBV) encoded latent membrane protein-1 (LMP-1) 30 bp deletion mutation effect on immune checkpoint regulation in EBV-associated Nasopharyngeal Carcinoma (NPC) and Non-NPC cancers

¹Naveed Ahmed, ^{2,3}Rohimah Mohamud, ^{1,3}Siti Asma Hassan, ^{3,4}Norhafiza Mat Lazim, ^{3,5}Wan Fatimah Wan Sohaimi, ^{3,6}Kueh Yee Cheng and ^{1,3}Chan Yean Yean*

¹Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

²Department of Immunology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

³Hospital Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan Malaysia

⁴Department of Otorhinolaryngology-Head & Neck surgery, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian 16150, Kelantan, Malaysia

⁵Department of Nuclear Medicine, Radiotherapy and Oncology, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, Kubang Kerian, Kota Bharu 16150, Kelantan, Malaysia

⁶Biostatistics and Research Methodology Unit, Health Campus, Universiti Sains Malaysia, Kubang Kerian, Kota Bharu 16150, Kelantan, Malaysia

ABSTRACT

Background: Nasopharyngeal carcinoma (NPC), lymphoma, breast, and gastric cancers are Epstein-Barr virus (EBV)-associated cancers. Presently, late-stage cancer is hard to manage. Therefore, early detection and precision cancer treatment are very important for cancer management. The detection of EBV LMP1 30 bp deletion in the biopsy samples of NPC patients were previously studied and proven to have an association with the development of NPC, but many questions remain unresolved, particularly on the effects of this mutation on the regulation of immune checkpoint receptors on immune cells (particularly regulatory T cells; Tregs) in cancers other than NPC. Methods: The current study was conducted at the Department of Medical Microbiology and Parasitology, School of Medical Sciences, Universiti Sains Malaysia (Health campus), Malaysia. The EBV LMP1 30 bp deletion mutation variants from whole blood samples from 29 patients with NPC, 29 non-NPC and 29 Healthy controls (HC) were determined using Polymerase Chain Reaction (PCR). Furthermore, the expression levels of immune checkpoint receptors (CD3, CD4, CD25, CD127, FOXP3, LAG-3, TIM-3, and TIGIT) was determined on Tregs using flow-cytometry assay. Statistical analysis was performed using SPSS programme. Results: The majority of NPC patients were Malay, followed by Chinese ethnicity (89.29% and 10.34%, respectively). Most of the NPC patients were diagnosed with WHO type III (n = 17, 58.62%), followed by WHO type II (n = 11, 37.93%), and WHO type I (n = 1, 3.44%). The locally advanced NPC stages were found in the vast majority of NPC patients (stage IV and stage III). LMP1 30-bp deletion mutation variant was detected in 8/29 (27.58%) of NPC patients, while none of the non-NPC and healthy control samples were found positive for LMP1 30 bp deletion mutation variant. The expression level of TIM3 on Tregs was 3.94 ± 5.79 in NPC, 17.2 ± 20.3 in non-NPC and 0.079 ± 0.25 in HC. For LAG3 the expression was 4.38 ± 9.25 in NPC, 1.83 ± 4.35 in non-NPC and 0.048 ± 0.18 in HC. For TIGIT the expression level was 29.5 ± 13.5 in NPC, 36.6 ± 22.0 in non-NPC and 1.03 ± 1.36 in HC. Conclusion: The results of this study contribute significantly to the understanding of EBV LMP1 30 bp deletion on the expression levels of immune checkpoint receptors in EBV-associated cancer patients which could relevant in identifying potential immune checkpoint inhibitor drugs for immunotherapy as well as prognostic biomarker in monitoring immunotherapy in EBV-associated (NPC and non-NPC) cancer patients.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

High Throughput DIA-NN Proteomics Identifies Altered Extracellular Protein Network in CRISPR-Targeted AGR2 Of Breast Cancer Cells.

¹Syazalina Zahari, ¹Saiful Effendi Syafruddin, ¹Low Teck Yew, ²Tiannan Guo, ¹M. Aiman Mohtar*

¹*UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, 56000, Kuala Lumpur, Malaysia.*

²*Guomics Laboratory of Proteomic Big Data, School of Life Science, Westlake University, 18 Shilongshan Street, Xihu District, Hangzhou, Zhejiang Province, China.*

ABSTRACT

Anterior Protein Gradient-2 (AGR2), a member of the protein disulphide isomerase (PDI) family located in the endoplasmic reticulum (ER), plays a pivotal role in catalyzing protein folding and thiol-disulphide interchange reactions. Its significance emerges from its overexpression in various epithelial-origin cancer types, accompanied by extracellular secretion, yet the functions of extracellular AGR2 (eAGR2) remain to be elucidated. This study dives into the intricate of eAGR2 functions, using breast cancer cells as a model. Employing CRISPR/Cas9 genome editing tool, AGR2 was targeted in three breast cancer cell lines: MCF-7, T47D, and 1833-BoM. Cell-based in-vitro assays of the AGR2-targeted cells confirmed AGR2's oncogenic nature in breast cancer. AGR2-repressed cells demonstrated reduced proliferative capacity, a hallmark of tumorigenesis. Furthermore, the cells displayed heightened sensitivity to conventional breast cancer drugs like doxorubicin, suggesting AGR2's involvement in drug resistance mechanisms. We also constructed four different structural AGR2 variants to modulate AGR2 secretion and found that the ER-retention motif might has a role in AGR2 secretion. To gain quantitative insights into eAGR2's role, state-of-the-art mass spectrometry Data Independent Acquisition-Neural Network (DIA-NN) methodology, were utilized. After subjecting the AGR2-null cells to serum starvation and secretome enrichment, DIA-NN mass spectrometry identified over 900 differentially expressed secretome proteins, with more than 300 consistently downregulated in all AGR2-targeted cell lines. Functional enrichment highlighted the involvement of these genes in critical cellular processes such as receptor signaling, secretion, extracellular vesicle biogenesis, cell adhesion, and inflammatory response. Protein-protein interaction network analysis uncovered a compelling interconnected hub genes within the downregulated proteins, shedding light on potential key players in the AGR2 signaling axis. This study unveiled bona-fide AGR2-modulated secretome that may influence the tumor microenvironment and breast cancer progression, thereby expanding the biology of eAGR2 function in the context of breast cancer.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Lipidome Analysis and Potential Lipid Biomarkers in Glioblastoma Tissue using Untargeted Lipidomic Analysis

¹Rashid K, ²Ibrahim K, ¹Wong JHD, ¹Ramli N

¹*Department of Biomedical Imaging, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia*

²*Department of Biomedical Science, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia.*

ABSTRACT

Background: Glioblastoma (GBM) is the lethal subtype of glioma, with a poor prognosis and patient survival rate. Dysregulation of the metabolism, particularly lipid homeostasis is linked to the GBM's aggressiveness and treatment resistance. This study aimed to characterise the lipidome composition and further investigate the potential lipid biomarkers in tissue samples of GBM patients. **Methods:** Human tumoural tissue lipidomes from 8 controls and 22 patients with glioblastoma were analysed using triple quadrupole liquid chromatography electrospray ionisation mass spectrometry. Differences in lipid profiles and differential lipid expression between GBM and controls were compared and investigated using univariate and multivariate statistical analysis. Receiver operating characteristic (ROC) analysis was performed, where the area under the curve (AUC) values larger than 0.75 were identified as potential lipid biomarkers. **Results:** Lipidome analysis revealed increased lipid composition with 7888 lipid compounds detected in GBM tissue. Several lipid subclasses including phosphatidylcholine, phosphatidylethanolamine, triacylglycerol, and fatty acyl were reduced, whereas ceramide was elevated in GBM tissue. Partial least squares-discriminant analysis (PLSDA) and hierarchical clustering analysis (HCA) demonstrated that GBM patient and control populations were visually separated by identified lipid biomarkers. Random forest analyses further predict the accuracy of 25 significant lipid species in GBM. Lipid metabolites, Cer 13:0;20/26:5 (AUC=1.00), DG 43:11 (AUC=0.94) and PC O-9:0_24:5 (AUC=0.89) were among the best lipids biomarker in GBM tissue. **Conclusion:** This study demonstrates lipid metabolic rewiring in glioblastoma, which may provide insight into its underlying molecular plasticity and progression for clinical benefits.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

TCR profiling of cytotoxic T cells targeting neoantigens from Colorectal Cancer Patients.

¹Nurul Ainaa' Adilah Rus Bakaruraini, ¹Ryia Illani Mohd Yunos, ²Isa Md Rose, ³Ismail Sagap, ¹Nurul Syakima Ab Mutalib, ¹Rahman Jamal, ¹Nadiyah Abu*.

¹UKM Medical Molecular Biology Institute (UMBI), UKM Medical Centre, Kuala Lumpur, Malaysia; ²Department of Pathology, UKM Medical Centre, Kuala Lumpur, Malaysia; ³Department of Surgery, UKM Medical Centre, Kuala Lumpur, Malaysia.

Corresponding author's email: nadiyah.abu@ppukm.ukm.edu.my

ABSTRACT

Background: Neoantigens, derived from non-synonymous mutations in a patient's genome could potentially become an ideal target for robust activation of anti-cancer response. Neoantigens are only expressed in tumor cells and are completely absent in normal cells. As such, neoantigens can become a precise target for cytotoxic T cells, especially in the application of cancer vaccines and adoptive T cell therapy (ACT). Method: PCR sequence-specific primer was done to check for the HLA status of ten high TMB CRC samples. Prediction of neoantigen was done using MuPeXi. Gene expression and immunogenicity analysis were done using the Class I immunogenicity tool. The affinity binding of neoantigen candidates was determined using NetMHC4.0. Primary cell culture work was done by isolating PBMCs from whole blood. Monocytes were isolated using the plastic adherence method and were cultured with maturation cytokines to form matured dendritic cells (mDCs). Validation of cellular markers i.e CD14, C83, CD86, CD3 and CD8 was done using FACS Verse. Ten μ M of the peptide was fed into mDCs before being co-cultured with CD8 T cells. Finally, neoantigen-reactive T cells were isolated using peptide dextramer and FACS Aria III. Total RNA was obtained and immunosequencing was performed. Briefly, 5' RACE was performed for reverse transcription and the products were amplified by PCR with corresponding specific primers. The libraries were sequenced using Illumina Novaseq instrument. Result: 5 samples were identified as HLA-A*2402 positive from 10 CRC samples. Data from combined analysis of MuPeXi, gene expression microarray, immunogenicity tool and NetMHC4.0 revealed one candidate that strongly binds to HLA-A*2402. Cytometric analysis revealed more than 90% of mDCs. Isolation of T cells using negative magnetic beads separation showed 93% of the T cell population. The sorting of T cells pulled down around 6% of neoantigen-specific T cells. Immunosequencing analysis of TCR revealed several dominant CDR3 clonotypes of alpha and beta chains in both CD8 and dextramer samples. Interestingly, the most dominant CDR3 sequence was also found in the tumor microenvironment, suggesting the presence of neoantigen-specific T cells from tumor-infiltrating lymphocytes. Conclusion: TCR sequence targeting specific neoantigen can be identified effectively using immunosequencing of sorted T cells by peptide dextramer. We provided a fast and effective method to culture and isolate neoantigen-specific T cells. In the future, the sequence of TCR can be cloned into healthy T cells for further functional validations against the peptide neoantigen.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Lutein inhibits the triple negative breast cancer cell progression by suppressing the protumour effects of tumour-associated macrophages.

¹Redzyque Ramza Ramli, ²Maryam Azlan, ²Nurul Asma Abdullah, ¹Siti Norasikin Mohd Nafi
and ¹Agustine Nengsih Fauzi

¹*Department of Chemical Pathology, School of Medical Sciences, Universiti Sains Malaysia Health Campus, 16150 Kota Bharu, Kelantan, Malaysia.*

²*School of Health Sciences, Universiti Sains Malaysia Health Campus, 16150 Kota Bharu, Kelantan, Malaysia,*

ABSTRACT

Tumour-associated macrophages (TAMs) play an important role in tumour development by modulating the M2-type macrophage (M2) polarisation in tumour microenvironment. Several phytochemicals have been reported to have inhibitory effects on M2 macrophage polarisation. Lutein (non-vitamin A carotenoids) is a natural compound with well-known for its medicinal benefits and has been shown to have promising anticancer properties. In the current study, we used in vitro assays to determine whether lutein has inhibitory effects on the M2 polarisation and its protumour functions towards MDA-MB-231 (triple negative breast cancer) cells. CD14⁺ isolation from the peripheral blood mononuclear cells (PBMCs) of healthy donors was performed using immunomagnetic beads to isolate monocytes and the monocytes were cultured with GM-CSF/LPS/IFN- γ or M-CSF/IL-4 to differentiate the monocytes into macrophages. MDA-MB-231 cell lines were co-cultured with human monocyte-derived macrophages (HMDMs) and subjected to the cell proliferation and transwell migration assays as well as the determination of cell surface markers. Our results showed that M2 macrophages accelerated the proliferation and migration of MDA-MB-231 cells and these protumour effects were inhibited by lutein. Additionally, lutein also significantly reduced the expression of M2 surface markers (CD206), and thus reversed the earlier elevation of CD206 induced by IL-4. Our data collectively imply that lutein may be an effective anticancer therapy or adjuvant for patients with TNBC.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Super Enhancer Upstream Regulators Drive KLF6-PDGFB Expression and mTORC1 Pathway Activation in Renal Carcinoma

¹Nurul Nadia Mohamad Zamberi, ²Jie Sheng Aik, ³Zulkifli Bin Md. Zainuddin, ¹M. Aiman Mohtar & ¹Saiful Effendi Syafruddin*.

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, Bandar Tun Razak, Kuala Lumpur 56000, Malaysia; ²St Edmund's College, University of Cambridge, Cambridge, CB3 0BN, United Kingdom; ³Department of Surgery, Level 8, Clinical Block, Hospital Canselor Tuanku Muhriz, Jalan Yaacob Latiff, Bandar Tun Razak, Kuala Lumpur 56000, Malaysia.

ABSTRACT

Kruppel-like factor 6 (KLF6), a super enhancer-driven transcription factor, supports clear cell renal cell carcinoma (ccRCC) growth by controlling cellular loop that links several ccRCC hallmark features including mTORC1 pathway hyperactivation. We have previously demonstrated that KLF6 transcriptionally regulates PDGFB expression, which will subsequently activate mTORC1 pathway in ccRCC via autocrine and paracrine manners. Nonetheless, whether the KLF6-PDGFB axis-mediated mTORC1 activation in ccRCC is also dependent on the function of super enhancer regulators, namely BRD4 and P300, remains elusive. Hence, to address this, we chemically perturbed BRD4 and P300 activities in a panel of ccRCC in-vitro models using inhibitor JQ1 and A485, respectively. This was followed by assessing (i) H3K27ac level, marker for active enhancer, via ChIP-qPCR, (ii) KLF6 and PDGFB expression level via qPCR and (iii) mTORC1 activity via Western blot. We found that BRD4 and P300 inhibitions reduced H3K27ac signals at several KLF6 locus constituent enhancers, in line with the critical roles of these factors in maintaining the super enhancer landscape. Moreover, the inhibition of BRD4 and p300 significantly reduced the expression of KLF6 and PDGFB in ccRCC cells, which consequently impaired the mTORC1 activity in the JQ1 and A485-treated cells. Our findings unravel the functional link between super enhancer upstream regulators, BRD4 and P300, and mTORC1 activation in ccRCC, which is via the transcriptional regulation of KLF6-PDGFB axis. Collectively, this study provides another important jigsaw for the construction of transcriptional dependency landscape in ccRCC, contributing to the holistic understanding on ccRCC pathogenesis for the development of novel and efficient clinical strategies

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Predictive Modelling of Cardiovascular Risk in Type 2 Diabetes Mellitus Patients: A K Nearest Neighbours Approach

Kee Ooi Ting¹, Harmiza Harun¹, Norlaila Mustafa², Nor Azian Abdul Murad¹, Chin Siok Fong¹, Rosmina Jaafar³, Hamat Hamdi Che Hassan², Mohd Zubir Suboh⁴, Noraidatulakma Abdullah^{1,5*}.

¹*Institut Biologi Perubatan Molekul (UMBI), Universiti Kebangsaan Malaysia (UKM), 56000 Kuala Lumpur, Malaysia.*

²*Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia (UKM), 56000 Kuala Lumpur, Malaysia*

³*Faculty of Engineering and Build Environment, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia*

⁴*Medical Engineering Technology Section, British Malaysian Institute, Universiti Kuala Lumpur, 53100 Kuala Lumpur, Malaysia*

⁵*Faculty of HealthSciences, Universiti Kebangsaan Malaysia (UKM), 50300 Kuala Lumpur, Malaysia*

**Correspondence: noraidatulakma.abdullah@ppukm.ukm.edu.my; Tel.: +60-3-9145-9072*

ABSTRACT

Background: Type 2 diabetes mellitus patients have a relatively higher risk of developing cardiovascular disease compared to non-diabetic individuals due to other microvascular complications. A prediction model is necessary to reduce the risk of cardiovascular disease development and the economic burden due to the treatment costs. Methods: This study has developed a cardiovascular disease risk prediction model with 220 participants recruited from Endocrine and Cardiology Clinic in Hospital Canselor Tuanku Muhriz (HCTM). After ranking the 15 features using ANOVA, several classifiers were trained with 80:20 data split and 10-fold cross validation using the organized data. Results: A K Nearest Neighbours model with 88.6% accuracy, 88.1% F1 score, and AUC of 0.9383 was selected. A graphical user interface was also developed to provide a user-friendly platform to run the prediction model. The prediction model developed in this study has a high accuracy and precision even though it is trained with a relatively small data size. Conclusion: Prediction models using machine learning should be introduced to the medical settings to assist in the development of cardiovascular complications among diabetes patients.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Alteration of Gut Microbiota in Systemic Lupus Erythematosus (SLE)

¹Qin Shi Cham*, ²Siok-Fong Chin, ¹Lydia Kamaruzaman, ¹Ruslinda Mustafar, ³Syahrul Sazliyana Shaharir, ⁴Asrul Abdul Wahab, ¹Rozita Mohd

¹UKM Medical Centre, Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia

²UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia

³Rheumatology Unit, Medical Department, Faculty of Medicine, Universiti Kebangsaan Malaysia

⁴Microbiology Department, Faculty of Medicine, Universiti Kebangsaan Malaysia

ABSTRACT

Alteration of gut microbiota could potentially be the initial step inducing dysregulation of the immune system in the gut-kidney axis that led to systemic inflammation. It is increasingly recognized in many diseases but studies in systemic lupus erythematosus (SLE) are still limited. This research was to determine the gut microbiota profile in SLE and its association with disease activity. A prospective study of 15 active renal SLE (aSLE) with age and sex matched control consisted of 15 renal SLE in remission (rSLE) and 15 non-renal SLE (nSLE) from PPUKM. Demographic data, stool samples, routine bloods and urine tests were collected and analysed. Stool DNA was extracted and 16S rRNA metagenomic sequencing was performed on an Illumina MiSeq Platform. The aSLE group had significant high SLEDAI-2K($p < 0.001$), low serum albumin($p < 0.001$) and total protein($p < 0.001$), lower serum haemoglobin($p = 0.031$) and high urea($p = 0.008$). No significant difference was observed in the richness and diversity (α -diversity), and the compositional difference (β -diversity) of bacterial species among the groups. However, the α -diversity metrics Simpson was significantly higher in rSLE than aSLE($p = 0.036$) which represent lower evenness of gut microbiome. The Firmicutes/Bacteroidetes ratio showed a decreasing trend between groups with aSLE(27.37 ± 80.31), rSLE(7.535 ± 12.23) and nSLE(5.314 ± 6.028), despite not significant. The most dominant phyla observed in aSLE was the proteobacteria, whereas actinobacteriota in the nSLE. Interestingly, LEfSe analysis revealed that bacteria genus Veillonella was enriched in the aSLE as compared to rSLE and nSLE, whilst Akkermansia was over-represented in the rSLE as compared to the nSLE. The genus Veillonella was positively correlated with disease activity ($r = 0.195$, $p = 0.2$) whereas Ruminococcus torques were negatively correlated with disease activity ($r = -0.166$, $p = 0.276$). The genus Veillonella(AUC=0.721, 95%CI 0.568-0.874) and Ruminococcus torques(AUC=0.694, 95%CI 0.536-0.853) were identified as a potential predictor for disease activity via ROC analysis.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Gene expression profiling of the extracellular vesicles from the insulin-resistant adipocytes.

¹Vigneswarry Dorairaj, ¹Nadiah Abu, ¹Nor Azian Binti Abdul Murad, ¹Rahman Jamal, ²Razman Jarmin, ¹Siti Aishah Sulaiman*

¹*UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia (UKM),
Jalan Yaacob Latif, Cheras, Kuala Lumpur, Malaysia.*

²*Department of Surgery, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia.*

*Corresponding author: sitiaishahsulaiman@ukm.edu.my

ABSTRACT

Background: Metabolic dysfunction-associated fatty liver disease (MAFLD) is the most common liver disease, often associated with obesity, insulin resistance/diabetes, or metabolic syndrome. Extracellular vesicles (EVs) are nano-sized particles secreted from the cells, with their cargos containing nucleic acids and proteins to mediate the communication between adipocytes and hepatocytes during MAFLD disease development. This study aimed to characterise the EVs isolated from insulin-resistant adipocytes and their cargos, focusing on gene expression profiles. **Methods:** Insulin-resistant (IR) and insulin-sensitive (IS) mature adipocytes were established in vitro. Their EVs were characterised via surface marker flow cytometry and morphology analysis using the transmission electron microscope (TEM) and dynamic light scattering (DLS). Total RNAs were isolated from the EVs and analyzed for differential gene expression using the human gene expression microarray (Agilent SurePrint v3). Significant genes were filtered based on the adjusted p-value (p_{adj}) < 0.05 and $-2 < \text{fold change} > 2$. Biological pathway and gene network analyses were performed using the g:profiler, STRING, and miRNet database. **Results:** EVs from IR and IS-adipocytes have similar surface markers (CD63, CD81, C9, and ALIX). Morphologically, both EVs were cup-shaped, though the EVs from IS-adipocytes were bigger in diameter (363.1 nm) than IR adipocytes (102.9 nm). The gene expression analysis identified that one gene was up-regulated, and 200 genes were down-regulated in EVs from IR-adipocytes compared to IS-adipocytes. Seventy-three genes were associated with EVs ($p_{adj}=1.1e-23$), and 75 genes were associated with adipocyte cell types ($p_{adj}=1.48e-2$). The top three biological pathways enriched are oxidative phosphorylation ($p_{adj}=3.88e-8$), reactive oxygen species (ROS) ($p_{adj}=6.09e-6$), and MAFLD ($p_{adj}=8.76e-4$). Network analysis identified one critical molecular network comprising the miR-146a-5p and its target genes (PTPN11, SLC25A6, and NDUFB2) involved in ROS regulation. **Conclusion:** Identifying the molecules inside the EVs from the adipocytes may provide a better understanding of the molecular communication between the adipocytes and hepatocytes, possibly leading to MAFLD development.

Keywords: Adipocytes, insulin resistance, extracellular vesicles, cell-cell communication, MAFLD

Acknowledgment: This study is sponsored by the Fundamental Research Grant Scheme (FRGS) from the Ministry of Higher Education (FRGS/1/2019/SKK08/UKM/03/8).

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Functional impact of selected DNA repair gene polymorphisms: insights from in silico analyses

Hilary Sito, Shing Cheng Tan*

UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Cheras, Kuala Lumpur, Malaysia

ABSTRACT

Background: DNA repair genes play a central role in maintaining genomic integrity. While single nucleotide polymorphisms (SNPs) in DNA repair genes are frequently associated with the risk of various diseases, the precise functional basis of these associations remains elusive. In this study, we investigated the functional implications of selected DNA repair gene polymorphisms using bioinformatics tools. Methods: Seven SNPs were selected for the in silico studies: ERCC1 rs3212986, ERCC2 rs1052555, ERCC2 rs1799793, XPC rs77907221, XRCC1 rs25487, XRCC1 rs1799782, and XRCC3 rs861539. A total of 32 different tools were used to predict the structure, domain, missense pathogenicity, splicing alteration, protein interactions, transcriptional and post-translational modification regulation, and the clinical effects of the SNPs. Results: ERCC2 rs1799793, XRCC1 rs25487, XRCC1 rs1799782, and XRCC3 rs861539 are missense SNPs, while ERCC1 rs3212986, ERCC2 rs1052555, and XPC rs77907221 are 3'-UTR, synonymous, and indel mutations, respectively. XRCC1 rs1799782 is predicted to have the highest probability of damaging effects, while ERCC2 rs1799793 is likely to be deleterious. Structural changes in XRCC1 rs1799782 and XRCC3 rs861539 may affect protein-protein interactions and downstream biological processes. XRCC3 rs861539 results in a loss of a phosphorylation site, impacting transcriptional regulation. ERCC1 rs3212986, ERCC2 rs1052555, and XPC rs77907221 were predicted to have altered splicing effects, with ERCC2 rs1799793 and XRCC1 rs25487 showing the most significant effects. Conclusion: Bioinformatics predictions of the functional effects, interactions and altered splicing effects of SNPs provide early insights into the underlying mechanisms of these SNPs in diseases, providing a basis for future experimental validation and better personalised treatment approaches.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Characterisation of extracellular vesicle released from hepatocytes with excessive lipid accumulation

¹Muhammad Nafiz Haidi Adrus, ¹Vicneswarry Dorairaj, ¹Nadiah Abu, ¹Nurul Syakima Ab Mutalib, ²Khairul Najmi Muhammad Nawawi, ¹Siti Aishah Sulaiman*

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia (UKM), Jalan Yaacob Latif, 56000 Cheras, Kuala Lumpur, Malaysia, ²Gastroenterology Unit, Department of Medicine, UKM Medical Centre, Jalan Yaacob Latif, 56000 Cheras, Kuala Lumpur, Malaysia.

ABSTRACT

Background: In metabolic dysfunction-associated fatty liver disease (MAFLD), fibrosis occurs as a result of uncontrolled wound-healing processes involving various interactions between several cell types, including hepatocytes, hepatic stellate cells (HSCs), Kupffer cells, as well as infiltrating immune cells. Studies suggested that extracellular vesicle (EV) released from hepatocytes following excessive lipid accumulation (lipotoxic) can contribute to cell interactions via their bioactive cargos during this event. One such is the miR-21, which has been reported to increase during fibrosis. This study aimed to characterise the EV from lipotoxic hepatocytes and its miR-21 expression. Methods: Human hepatocyte cell line (HepG2) treated with 0.5mM palmitate (PA) conjugated with 5% bovine serum albumin (BSA) (lipotoxic), and HepG2 cells treated with 5% BSA (CON) grown in serum-free media for 48 hours. Lipid accumulation was confirmed using the Oil Red O (ORO) staining and lipid marker expression. EVs were isolated and characterised using Transmission Electron Microscopy (TEM), Dynamic Light Scattering (DLS), and flow cytometry to assess their morphology, size, and surface markers. Total RNA was isolated from the EVs, and expression of miR-21 was measured using the Real-Time quantitative PCR method. Results: Lipotoxic hepatocytes have higher lipid accumulations ($P < 0.001$) with significant upregulation of Fatty acid synthase (FAS) gene expression ($P < 0.05$) compared to CON. EVs from lipotoxic and CON hepatocytes have similar surface markers, such as CD63, CD81, and CD9. Morphologically, both EVs were cup-shaped with similar sizes. Within the EV from lipotoxic hepatocytes, there was an enrichment of miR-21 expression (59.4-fold, P -value < 0.0001). Conclusion: EV miR-21 enrichment in lipotoxic hepatocytes may suggest that miR-21 could mediate fibrosis when the surrounding liver cells, such as HSC and Kupffer cells, take in these EVs.

Keywords: Lipid accumulation, hepatocytes, extracellular vesicle, cell-cell communication, MAFLD, miR-21.

Acknowledgment: This study is sponsored by the UKM University Grant (GUP-2021-067).

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Metabolic Signatures in Obesity: Discovery of Promising Independent Biomarkers

^{1,2}Anis Adibah Osman, ^{1,3}Noraidatulakma Abdullah, ¹Siok-Fong Chin, ⁴Lay-Kek Teh, ¹Nor Azian Abdul Murad, ¹Rahman Jamal*

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, Bandar Tun Razak, 56000 Cheras, Wilayah Persekutuan, Kuala Lumpur, Malaysia; ²Training Division, Ministry of Health, 62675 Putrajaya, Malaysia; ³Faculty Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia; ⁴Integrative Pharmacogenomics Institute (iPROMISE), Faculty of Pharmacy, UiTM Puncak Alam Campus, Bandar Puncak Alam, 42300 Puncak Alam, Selangor

ABSTRACT

Obesity is characterized by the excessive accumulation of body fat. This condition will lead to changes in metabolic profile which reflects the alteration of key metabolites and the biochemical pathways. Therefore, the discovery of signatory metabolites in obese individuals will help us to elucidate the mechanisms and pathogenesis of obesity. In this nested case-control study, a serum metabolomics profiling was performed of 60 samples from The Malaysian Cohort comprising of lean and obese participants, by utilizing BMI values of 18.5-22.9 kg/m² and ≥ 30 kg/m², respectively. A total of 85 significant metabolic features were identified between cases and controls. Due to weak correlations among metabolites detected, a metabolite model was built using a stepwise logistic regression and this model revealed 6 highly promising markers that are linked to obesity, and this showed performance of AUC = 0.950. Metabolites with elevated risk (OR > 1) were suggested to exhibit impairment in lipid metabolism. In order to assess the individual contribution of each metabolite to predicting obesity, we conducted multiple logistic regression analysis. All metabolites displayed enhanced capabilities in predicting obesity following the adjustment for confounding variables. 14-methylheptadecanoic acid (OR = 3.952; 95% CI = 1.396-11.184), 4'-apo-beta,psi-caroten-4'-al (OR = 1.180; 95% CI = 1.062-1.311) and 7,8-Dihydro-3b,6a-dihydroxy-alpha-ionol 9-[apiosyl-(1->6)-glucoside] (OR = 0.873; 95% CI = 0.765-0.995) exhibited independent associations after being adjusted for age, gender, race, occupation, and smoking status. This metabolite prediction could provide a clearer understanding of the development of obesity and potentially facilitate personalized medical approaches.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

POSTER

P1

Expression of Receptor Activator of Nuclear Factor-Kappa B (RANK) Protein in ER-Positive Pre-Therapeutic Breast Cancer Tissues

Alaa Siddig, Department of Pathology, School of Medical Sciences, Universiti Sains Malaysia

P2

The Incidence Rate and Characteristics of Positive Individuals for COVID-19 in Perak, Malaysia: July – December 2021

Filza Noor Asari, Institute for Public Health, National Institute of Health, Ministry of Health

P3

Clinical Validation of a CircANAPC7 Basescope Probe for the Detection of FOLFOX Chemoresistant Colorectal Cancer Via In Situ Hybridization

Nur Syafiqah Abd Hakim, UKM Medical Molecular Biology Institute (UMBI)

P4

Assessment of WGS In Silico HLA Typing Tools Against Molecular HLA Typing Method

Hamidah Akmal Hisham, Molecular Pathology Unit, Institute for Medical Research, National Institute of Health

P5

Microbiome, Race and Breast Cancer: Is There an Asian-Specific Gut Microbiome That May Affect Breast Cancer?

Audrey Weng Yan Lee, Cancer Research Malaysia

P6

Navigating Cardiovascular Risk in Type 2 Diabetes: Potential Roles of Lipid Peroxidation and Cardiometabolic Markers

Harmiza Harun, UKM Medical Molecular Biology Institute (UMBI)

P7

The Effect of ZFAS1 Silencing on Cell Viability in Glioblastoma Cell Lines LN18 (PTEN Wildtype) and U87MG (PTEN Mutant)

Engku Maryam Afifah Engku Omar Amiruddin, Institute of Medical Molecular Biotechnology (IMMB), Faculty of Medicine, Universiti Teknologi MARA

P8

Hyperglycemic Induced Pluripotent Stem Cell (iPSC) Reprogramming: Unveiling the Bittersweet Epigenetic Memories in iPSCs and Their Differentiated Endothelial Cells

Khoo Tze Sean, UKM Medical Molecular Biology Institute (UMBI)

P9

Hsa-miR-181a-5p, Hsa-miR-182-5p, and Hsa-miR-26a-5p As New Biomarkers for Undetermined BCR-ABL1 in Chronic Myeloid Leukemia Patients Treated With Tyrosine Kinase Inhibitors

Aliza Mohd Yacob, Cancer Research Centre, Institute for Medical Research, National Institutes of Health

P10

Polygenic Risk Score Significantly Predicted Coronary Heart Disease Death Among Those With Type 2 Diabetes in The Malaysian Cohort

Khairul Anwar Zarkasi, UKM Medical Molecular Biology Institute (UMBI)

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

P11

Revealing Uncharted Depths of the Dark Proteome: A Re-examination of Mass Spectrometry Data Uncovers New Insights into Colorectal Cancer Microproteins

Low Teck Yew, UKM Medical Molecular Biology Institute (UMBI)

P12

Demethylbelamcandaquinone B from *Marantodes pumilum* var. *alata* (Blume) Kuntze Inhibits Osteoclast Differentiation in RAW264.7 Cells

Haryati Ahmad Hairi, Department of Biochemistry, Faculty of Medicine, Manipal University College

P13

GPC3 Alterations and Interactions in Lung Adenocarcinomas: A Comprehensive Data Mining Study of Publicly Available Datasets

Tan Shing Cheng, UKM Medical Molecular Biology Institute (UMBI)

P14

Phytochemical Profiling and DPPH (2,2-diphenyl-1-picrylhydrazyl) Radical Scavenging Activities of the Crude Extract Ethanol *Laportea decumana* Rob. Wedd

Lukman La Basy, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia

P15

Unraveling AGR2 Regulation in Breast Cancer: Insights from RNA-Seq Analysis of CRISPR/Cas9 Mediated Targeting

M. Aiman Mohtar, UKM Medical Molecular Biology Institute (UMBI)

P16

Integrative Pipeline to Analyze Immune Cells in Digital Pathology Images

Haslina Makmur, Cancer Research Malaysia

P17

KCNN4 Mutation Presenting With Non-Immune Hydrops Haemolytic Anaemia

Nur Mashitah Abdullah, Department of Pediatrics, Hospital Pakar Kanak-Kanak (HPKK), The National University of Malaysia

P18

Risk Factors of Long COVID in Klang Valley Malaysia: A Snapshot from the COVGEN Study

Noraidatulakma Abdullah, UKM Medical Molecular Biology Institute (UMBI)

P19

Development of Breast Cancer Spheroids as a Model for a Pre-Clinical Evaluation Using Vitamin C As Anti-Cancer Agent

Ali Saeed Ali Mussa, Department of Haematology, School of Medical Sciences, Universiti Sains Malaysia

P20

Dipeptidyl Peptidase 4 (DPP IV) Inhibitory Activity of Leaf Extract and Sesquiterpene Lactones Isolated From *Smallanthus sonchifolius*

Hady Anshory Tamhid, Faculty of Pharmacy, Universitas Gadjah Mada, Indonesia

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

P21

A KLF6-Dependent PDGFB-mTOR Signalling Loop Links Lipid Homeostasis and Tumor Growth in Renal Carcinoma

Saiful Effendi Bin Syafruddin, UKM Medical Molecular Biology Institute (UMBI)

P22

Advancements in Amber Suppression Technology: A Window into Molecular Interactions of Human Viruses

Nur Firdaus Isa, Department of Biotechnology, Kulliyah of Science, International Islamic University Malaysia

P23

Metabolomic Fingerprinting and Footprinting Analyses on TNBC sh-TINCR Cell Lines

Mohd Ikhwani Ismail, Faculty of Pharmacy, Universiti Teknologi MARA

P24

Evaluation and Optimisation of Adipogenic Differentiation Protocol for the Human Adipocyte Tissue-derived Stem Cells

Siti Aishah Sulaiman, UKM Medical Molecular Biology Institute (UMBI)

P25

SLC6A14 Expression and Methionine Uptake Increased in Advanced Colorectal Cancer Cells Stage

Hazwani Mohd Yusof, Department of Biochemistry, Faculty of Medicine, Manipal University College Malaysia

P26

Optimization of Annealing Temperature for Dengue Virus Serotyping Using Multiplex Polymerase Chain Reaction (PCR) Technique

Nurul Syafiqah Nabilah Abdul Kadir, UKM Medical Molecular Biology Institute (UMBI)

P27

Current Status of Predictive Serum Biomarkers for Polyps and Colorectal Cancer

Maryam Fatimah Abu Bakar, Department of Public Health Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia

P28

High-Throughput Genotyping of Asian-Specific Genetic Variants Associated With Colorectal Cancer Risk in the Malaysian Population: A Pilot Study

Sharzehan Mohamad Ayub Khan, UKM Medical Molecular Biology Institute (UMBI)

P29

Novel Ryr2 Receptor Gene Mutation Presenting with Cardiac Tumour and Refractory Arrhythmias: Case Report and Systemic Review

Nur Mashitah Abdullah, Department of Pediatrics, Hospital Pakar Kanak-Kanak (HPKK), The National University of Malaysia

P30

Whole exome sequencing (WES) in Mullerian Agenesis and Anorectal Anomaly: A Case Study

Nor Azian Abdul Murad, UKM Medical Molecular Biology Institute (UMBI)

P31

Integrin Beta 1/CD29: A preliminary Investigation into a Possible GBM EV Surface Marker

Johannes Low Jun Wei, UKM Medical Molecular Biology Institute (UMBI)

P32

Radioactive Iodine Refractory Papillary Thyroid Cancer: ABC Transporter Genes Under the Microarray Lens

Nur Fadhina Mohamad Pakarulrazy, UKM Medical Molecular Biology Institute (UMBI)

P33

CovGen Study: Dissecting COVID-19 Symptom Severity and Risk Factors

Nor Adzimah Johdi, UKM Medical Molecular Biology Institute (UMBI)

P34

High Glucose Level Reduces Cisplatin Chemosensitivity in Cervical Cancer Cells

Siti Aisyah Md Sanib, Cluster of Integrative Physiology and Molecular Medicine, Universiti Kuala Lumpur
Royal College of Medicine

P35

Chemical Characterization, Antioxidant Potential and In Vitro Anticancer Properties of Rice (*Oryza sativa* var Bajong) from Sarawak

Lam Xiu Qian, Department of Paraclinical Sciences, Faculty of Medicine and Health Sciences,
Universiti Malaysia Sarawak

P36

LRG1 Induces Lysosomal Autophagy via Dysregulation of the mTOR Pathway

Lim Su Min, UKM Medical Molecular Biology Institute (UMBI)

P37

Chemosensitization of TINCR: Potential Key Regulator to Chemoresistance in Triple Negative Breast Cancers (TNBCs)

Ezanee Azlina Mohamad Hanif, UKM Medical Molecular Biology Institute (UMBI)

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Expression of Receptor Activator of Nuclear Factor-kappa B (RANK) Protein in ER-Positive Pre-Therapeutic Breast Cancer Tissues

¹Alaa Siddig, ¹Siti Norasikin Mohd Nafi, ²Sarina Sulong, ^{3,5}Tengku Ahmad Damitri Al-Astani Tengku Din, ^{4,5}Maya Mazuwin Yahya, ^{1,5}Wan Faiziah Wan Abdul Rahman*

¹Department of Pathology, ²Human Genome Centre, ³Department of Chemical Pathology, ⁴Department of Surgery, School of Medical Sciences, Universiti Sains Malaysia, Health Campus, 16150 Kubang Kerian, Kelantan, Malaysia. ⁵Breast Cancer Awareness and Research Unit, Hospital Universiti Sains Malaysia, 16150, Kubang Kerian, Kelantan, Malaysia

Correspondence: wfaiziah@usm.my

ABSTRACT

Background: RANK-RANKL signalling is a potential therapeutic target for breast cancer, regulating breast tumour initiation by modulating the proliferative response to progesterone hormone and mammary stem cells extension. RANK expression correlates with poor survival and chemotherapy resistance in estrogen receptor (ER)-negative breast cancer. This study explores the expression pattern of RANK in ER positive-breast cancer surgical tissues in relation to other clinicopathological features.

Method: This is a cross-sectional study conducted at Hospital University Science Malaysia (USM). The protein expression of RANK was assessed in (n=65) ER-positive breast cancer pre-therapeutic tissue samples using immunohistochemical (IHC) staining. The intensity and the percentage of stained tumour cells were used to determine the H-score. Results: 72.3% (47/65) of ER-positive breast cancer have higher expression of RANK protein (H-score > 200). Higher expression of RANK was significantly associated with negative lymph node invasion status (22/25) P <0.045. However, the RANK protein's expression level did not exhibit significant associations with other clinicopathological parameters. Conclusion: Intriguingly, the findings suggests that RANK protein might play an inhibitory role in ER-positive breast cancer, in contrast to its observed implications in ER-negative breast cancer. However, further research is crucial to unravel their complex role.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

**The incidence rate and characteristics of positive individuals for COVID-19 in Perak,
Malaysia: July – December 2021**

Filza Noor Asari^{1,*}, Eida Nurhadzira Muhammad¹, Muhammad Faiz Mohd Hisham¹, Norizah
Ismail², Husna Khairam², Mohd Emir Shafiq Suhaimi², Zahrul Laili Abd Hadi²

*Institute for Public Health, National Institute of Health, Ministry of Health¹ Hospital Raja Permaisuri
Bainun Ipoh²*

ABSTRACT

Background: As of 12th August 2023, Malaysia had more than 5.1 million COVID-19 cases and Perak reported 243,083 confirmed cases with 2,168 deaths. Between 1st July 2021 and 31st December 2021, a total of 45,329 COVID-19 samples were screened and tested by real-time RT-PCR. This study aimed to determine the incidence rate of COVID-19 infection based on districts in Perak and explore the characteristics of patients who tested positive for COVID-19. Methods: This is a retrospective study. Secondary data of Perak COVID-19 cases were obtained from Sistem Informasi Makmal Kesihatan Awam (SIMKA), a national reporting system by the Disease Control Division, Ministry of Health Malaysia. All 11 districts in Perak were included in this study. All COVID-19 positive cases confirmed by Real-time PCR between 1st July 2021 to 31st December 2021 were included in this study. All data were run for descriptive analysis using IBM SPSS v28. Results: A total of 9,432 cases were confirmed positive for COVID-19 infection. The overall incidence rate in Perak was 377 cases per 100,000 population. The majority of them were among Malay ethnicity (74.3%) and female contributed 53.8% of total cases. High cases were seen among the elderly aged >60 years and middle age group 30-39 years with the percentage of 21.1% and 20.3%, respectively. The incidence rate of COVID-19 was highest in Kuala Kangsar (605 cases per 100,000 population), followed by Hulu Perak (548 cases per 100,000 population), Larut Matang & Selama (512 cases per 100,000 population), Hilir Perak (485 cases per 100,000) and Manjung (341 cases per 100,000). Among the reasons for the COVID-19 screening were those categorized as a probable case (25.3%, n=2238), close contact (24.8%, n=2196), symptomatic screening (15.3%, n=1463), severe acute respiratory illness, SARI (9.1%, n=988), Pre-admission (4.4%, n=622) and others (21.2%, n=1,921). Conclusion: Data provide useful insights of COVID-19 incidence rate in Perak and their socio-demographic characteristic and an in-depth understanding of the infection trend. These data will be helpful in evaluating further infection rates and improve future protective measures for the Perak population.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Clinical Validation of a CircANAPC7 Basescope Probe for The Detection of FOLFOX Chemoresistant Colorectal Cancer Via In Situ Hybridization

1Nur Syafiqah Abd Hakim, ²Azyani Yahya, ³Marfu'ah Nik Eezamuddeen, ¹Nurul Aina Fatiha Jamli, ¹Puteri Hanis Nafisah Megat Kamil Azmi and ¹Nadiah Abu*

¹UKM Medical Molecular Biology Institute (UMBI), UKM Medical Centre, Kuala Lumpur; ²Department of Histopathology, Hospital Canselor Tuanku Muhriz UKM, Kuala Lumpur; ³Department of Oncology, Hospital Canselor Tuanku Muhriz UKM, Kuala Lumpur

ABSTRACT

Background: FOLFOX chemotherapy is one of the standard regimens for treating colorectal cancer (CRC). However, overcoming chemoresistance is a major hurdle in achieving favorable results. Hence, it is imperative to identify biomarkers that can predict the response to FOLFOX treatment, which can aid in better patient management. Circular RNAs (circRNAs) have emerged as promising biomarkers in cancer research due to their stability and dysregulation in various diseases. Our previous results have indicated that circANAPC7 is implicated in chemoresistance in CRC. Therefore, we aim to identify circANAPC7 using a Basescope probe method. Methods: In this study, we aimed to clinically validate a Basescope probe for detecting circANAPC7 expression in FOLFOX chemoresistant CRC via in situ hybridization. The patients were divided into two groups, responders and non-responders, based on their clinical response to FOLFOX treatment. This study analyzed tumor tissue samples from 26 CRC patients, which consisted of 14 non-responder and 12 responder patients who underwent FOLFOX treatment. Staining scores were assigned to quantify the expression levels of circANAPC7 in the samples. Results: We identified that non-responder samples exhibited higher total staining scores, indicating a higher circANAPC7 expression compared to responder samples. Conclusion: These findings validate the clinical utility of the CircANAPC7 Basescope probe as a potential biomarker in predicting FOLFOX treatment response in colorectal cancer patients, enabling personalized treatment decisions and improved management of chemoresistant cases.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor

Assessment of WGS in silico HLA typing tools against molecular HLA typing method

¹Hamidah Akmal Hisham*, ²Abdul Malik Ahmad, ²Akzam Saidin, ¹Tan Geok Wee, ³Tan Tee Yong, ³Wong Chun Ying and ¹Tan Lu Ping

¹*Molecular Pathology Unit, Institute for Medical Research, National Institute of Health, Malaysia.*

²*Novocraft Technologies Sdn Bhd, Petaling Jaya, Malaysia.*

³*Hospital Umum Sarawak, Kuching, Malaysia*

ABSTRACT

The HLA genes are a group of highly polymorphic genes that are in the MHC region on chromosome 6. The HLA genotype affects the presentability of tumour antigens to the immune system. Therefore, methods to accurately type HLA alleles are critical to study differences in immune response between cancer patients. PCR-based methods are the current gold standard, but large-scale datasets with PCR-based HLA genotypes are rarely available. A variety of methods for in silico NGS-based HLA genotyping have been developed, bypassing the need to determine these genotypes with separate experiments. However, there is currently no consensus on the best performing tool. We performed whole genome sequencing and molecular HLA typing for HLA-A and HLA-B, on 46 healthy individuals of Bidayuh ethnicity. We used novoHLA and T1K in silico HLA typing pipelines to perform HLA typing on WGS data and benchmarked against molecular HLA typing results. We aimed to evaluate the precision and accuracy of in silico HLA typing tools in identifying HLA-A and HLA-B. Our assessment showed that there is a high concordance between in silico HLA typing tools and molecular HLA typing. The results of in silico HLA typing pipelines performance assessment on Bidayuh WGS dataset will be presented.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Microbiome, Race and Breast Cancer: Is there an Asian-specific gut microbiome that may affect breast cancer?

¹Li-Fang Yeo, ¹Audrey Weng Yan Lee, ¹Seow Fong Chin, ¹Soo-Hwang Teo, ¹Jia Wern Pan*

#Co-first authors

¹Cancer Research Malaysia, Subang Jaya Medical Centre, Subang, 47500 Selangor, Malaysia

ABSTRACT

Race has long been considered a contributive factor affecting breast cancer, and in recent years, microbiome diversity. The efforts to identify race-specific microbiomes have grown in recent years with the idea of personalized medicine, yet such studies claiming race-specific microbiomes in health and disease continue to be divided. Thus, we conducted a systematic review to address the question: "Is there an Asian-specific gut microbiome that may affect breast cancer?" We conducted a literature search on PubMed, Embase, Cochrane, and Web of Science. Publications included were limited to original research studying breast cancer and gut microbiomes on Asian populations. At first glance, literature seems to suggest that there is a racial difference in the microbiome of breast cancer patients. However, there is a great lack of Asian representation in international breast cancer and microbiome studies with Caucasian cohorts outnumbering Asian cohorts by up to fifty times. In-depth investigation of these studies revealed gross overgeneralization of racial groups. Heterogenous sub-groups of Asian, Latin American, and mixed-race participants were categorized into inappropriate racial groups. Findings from such multi-racial studies in breast cancer were repeated widely across literature without scrutiny and eventually accepted as dogma. The impact of lifestyle factors, such as socioeconomic status and unique behaviors caused by diet and cultural differences, on microbiome and breast cancer profiles were repeatedly acknowledged but remain unaccounted for. Moving forward in studying the effect of a race-specific microbiome in breast cancer, our findings highlight that Asians should not be studied as a homogenous group as it may result in inaccurate conclusions affecting real-world applications such as precision medicine. We hope that an improved understanding of Asian cohorts may improve on the granularity of the effect of race on microbiomes and breast cancer.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor

Navigating Cardiovascular Risk in Type 2 Diabetes: Potential Roles of Lipid Peroxidation and Cardiometabolic Markers.

¹Harmiza Harun, ¹Kee Ooi Ting, ²Norlaila Mustafa, ¹Nor Azian Abdul Murad, ¹Chin Siok Fong,
³Rosmina Jaafar, ²Hamat Hamdi Che Hassan and ^{1,4}Noraidatulakma Abdullah*.

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia (UKM), 56000 Kuala Lumpur, Malaysia

²Department of Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia (UKM), 56000 Kuala Lumpur, Malaysia

³Faculty of Engineering and Built Environment, Universiti Kebangsaan Malaysia, Bangi, 43600, Malaysia

⁴Faculty of Health Sciences, Universiti Kebangsaan Malaysia (UKM), 50300 Kuala Lumpur, Malaysia

*Correspondence: noraidatulakma.abdullah@ppukm.ukm.edu.my; Tel.: +60-3-9145-9072

ABSTRACT

Type 2 diabetes poses an elevated risk of cardiovascular disease, contributing to higher morbidity and mortality rates. Identifying individuals at stake in the early stages of CVD is crucial for effective intervention. In this regard, lipid peroxidation and cardiometabolic markers offer insights into the intricate interplay between diabetes and cardiovascular risk, aiding in early detection and risk assessment. This study aims to synthesize evidence on lipid peroxidation and cardiometabolic markers associated with the early onset of cardiovascular disease, shedding light on their potential clinical utility. A systematic search was conducted in Pubmed, Web of Science and Scopus to identify relevant studies published between January 2017 and August 2022. We included studies investigating lipid peroxidation and cardiometabolic markers implicated in the early progression of cardiovascular disease among type 2 diabetes patients. This systematic review has been registered with PROSPERO: CRD42022356566. The search yielded 76 studies, of which seven that met the criteria were finally included. Through meticulous analysis, malondialdehyde was identified as the promising candidate for lipid peroxidation markers in type 2 diabetes with cardiovascular risk. Other relevant markers investigated include fluorescent damage products of lipid peroxidation, lipid peroxides, thiobarbituric acid reactive substances, oxidized-LDL, arylesterase PON1, and ferric-reducing ability of plasma. The selected studies were of good quality, presenting a median score of 18 (13-25) based on the Downs and Black risk assessment checklist. This systematic review highlights the role of lipid peroxidation and cardiometabolic markers in unveiling the early cardiovascular risk among type 2 diabetes individuals. These biomarkers hold the potential for early detection and risk stratification of cardiovascular disease as well as improving the management of cardiovascular complications in the vulnerable population. Further research is essential to validate and establish the clinical utility of these biomarkers within diverse type 2 diabetes populations.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

The Effect of ZFAS1 Silencing on Cell Viability in Glioblastoma Cell Lines LN18 (PTEN Wildtype) and U87MG (PTEN Mutant)

¹Engku Maryam Afifah Engku Omar Amiruddin, ^{1,2}Amirah Abdul Rahman*, ²Siti Hamimah Sheikh Abdul Kadir, ³Kamariah Ibrahim and ²Nora Julianna Osman

¹*Institute of Medical Molecular Biotechnology (IMMB), Faculty of Medicine, Cawangan Selangor, Kampus Sungai Buloh, Universiti Teknologi MARA (UiTM), 47000 Sungai Buloh, Selangor, Malaysia*

²*Faculty of Medicine, Universiti Teknologi MARA (UiTM), Cawangan Selangor, Kampus Sungai Buloh, 47000 Sungai Buloh, Selangor, Malaysia*

³*Department of Biomedical Science, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia.*

ABSTRACT

Recent research has revealed the multifaceted role of ZNF1 antisense RNA 1 (ZFAS1) in cancer, where it can function as either a tumor suppressor or an oncogene, exerting a critical influence on various cellular processes. Intriguingly, ZFAS1 exhibits upregulated expression in numerous cancers, such as colorectal and ovarian cancers, but is downregulated in breast cancer. Glioblastoma (GBM) demonstrates elevated ZFAS1 expression relative to normal brain cells. Nevertheless, emerging evidence suggests that the expression of ZFAS1 may vary based on cancer cell types and traits. This study aims to elucidate the impact of ZFAS1 silencing on GBM cancer cells, considering their PTEN (phosphatase and tensin homolog) status, focusing on cell viability and clonogenicity. Real-time quantitative PCR (RT-qPCR) was utilised to quantify ZFAS1 expression in Normal Human Astrocytes (NHA) and GBM cell lines (LN18 and U87MG) before siRNA transfection. si-ZFAS1 transfection efficiency was validated by RT-qPCR and siGLO fluorescence imaging. Cell viability was assessed using the MTT assay, and clonogenicity was evaluated. Despite the evidence on ZFAS1 being overexpressed in GBM relative to normal brain cells, our findings indicate that NHA cells exhibit higher ZFAS1 expression compared to LN18 and U87MG. Notably, LN18, a PTEN wild-type cell line, demonstrated significantly higher ZFAS1 expression than U87MG, a PTEN-mutated cell line. LN18 si-ZFAS1 silencing efficiency was 90.22%, and U87MG was 62.77%. Cell viability assays at 24-, 48-, and 72-hours post-siRNA transfection showed reduced LN18 cell viability (75.74%, 78.39%, and 59.68%) and moderate effects on U87MG (103.83%, 87.48%, and 92.96%) compared to non-targeting controls. In conclusion, ZFAS1 silencing significantly reduces cell viability, particularly in LN18 (PTEN wild-type), suggesting ZFAS1's oncogenic role in GBM development, with its impact influenced by GBM subtype or characteristics. Further investigation on the mechanism of action of ZFAS1 in GBM cells is warranted.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Hyperglycemic Induced Pluripotent Stem Cell (iPSC) Reprogramming: Unveiling the Bittersweet Epigenetic Memories in iPSCs and Their Differentiated Endothelial Cells.

¹Tze Sean Khoo, ¹Rahman Jamal, ²Nur Azurah Abdul Ghani, ³Hafiza Alauddin, ³Hamidah Hussin, ¹Nor Azian Abdul Murad*

¹ *UKM Medical Molecular Biology Institute, National University of Malaysia*

² *Obstetrics & Gynaecology Department, UKM Medical Centre, National University of Malaysia*

³ *Haematology Unit, Pathology Department, UKM Medical Centre, National University of Malaysia.*

ABSTRACT

Background: Induced pluripotent stem cell (iPSC) offers great promise in regenerative medicine due to its pluripotency and indefinite self-renewal capabilities. However, most if not all iPSC lines were reprogrammed and maintained under hyperglycemic conditions, which raises safety concerns. Despite hyperglycemia is known to be detrimental to adult cells, its impact on iPSC has not been well-studied. Hence, this study aimed to elucidate the impact of hyperglycemic iPSC reprogramming on the DNA methylation profile and functions of the resulting iPSCs and their differentiated endothelial cells (EC). **Methods:** Human umbilical vein endothelial cells were first isolated and expanded from a healthy donor in normoglycemic (LG-i-EC) or hyperglycemic (HG-i-EC) conditions. LG-i-EC and HG-i-EC were then reprogrammed into iPSC under normoglycemic (LG-iPSC) and hyperglycemic (HG-iPSC) conditions, respectively. Following successful reprogramming, HG-iPSC was expanded under normoglycemic condition (HGLG-iPSC) to elucidate the epigenetic memory retained after cessation of hyperglycemic exposure. iPSCs (LG-iPSC, HG-iPSC, and HGLG-iPSC) were then differentiated into endothelial cells (LG-iPSC-EC, HG-iPSC-EC, HGLG-iPSC-EC) in their respective conditions. All iPSCs and their differentiated ECs were characterized by flow cytometry, immunocytochemistry staining of cell-specific markers and were subjected to global DNA methylation microarray analysis and functional assessment. **Results:** All iPSCs expressed pluripotent markers SSEA-4, TRA-1-60, TRA-1-81 and OCT4, and proliferate at similar rates despite differences in glucose levels. However, differential methylation analysis revealed 2400 CpG sites differentially methylated in HGLG-iPSC compared to LG-iPSC, with over 70% of the 2400 epigenetic memories inherited by their differentiated ECs. Molecular pathway analysis implicated alterations in the transendothelial leukocyte migration pathway in HGLG-iPSC-EC. Functional assays demonstrated increased transendothelial monocyte migration activity in HGLG-iPSC-EC compared to LG-iPSC-EC, indicative of endothelial dysfunction. **Conclusion:** In conclusion, iPSC reprogramming in normoglycemic, not hyperglycemic, conditions is crucial, to avoid persistent epigenetic memories that lead to dysfunctional ECs, rendering them unsuitable for clinical applications in regenerative medicine.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Hsa-miR-181a-5p, hsa-miR-182-5p, and hsa-miR-26a-5p as new biomarkers for undetermined BCR-ABL1 in chronic myeloid leukemia patients treated with tyrosine kinase inhibitors

¹Aliza Mohd Yacob*, ²Nor Asiah Muhamad, ¹Hamidah Akmal Hisham, ³Latifah Ibrahim, ^{1,4}Shuhaila Mat Sharani, ⁵Erni Zuraida Ali, ¹Ivyna Pau Ni Bong, ¹Azli Ismail, ⁶Azura Mohd Noor and ¹Yuslina Mat Yusoff

¹Cancer Research Centre, Institute for Medical Research (IMR), National Institutes of Health (NIH), Selangor; ²Sector for Evidence Based Healthcare, NIH; ³Special Research Centre, IMR, NIH; ⁴Biomedicine Program, School of Health Science, Universiti Sains Malaysia (USM), Kelantan; ⁵Nutrition, Metabolic and Cardiovascular Research Centre, IMR, NIH; ⁶Infectious Disease Research Centre, IMR, NIH; Malaysia

ABSTRACT

Background: Chronic Myeloid Leukemia (CML) patients are treated with tyrosine kinase inhibitors (TKIs) and showed a high success rate. However, molecular relapse occurred 50% after discontinuation, with 80% occurring within the first six months. The molecular response is monitored and guided by BCR-ABL1 levels. The gold standard RT-qPCR measures gene expression (mRNA), and the new digital PCR has improved gene-level (DNA) sensitivity. Therefore, a complementary method is needed. In a previous study, we identified and validated three potential miRNAs (for translational repression / mRNA degradation) as therapeutic biomarkers for monitoring CML. Hsa-miR-181a-5p at below current limit of detection, hsa-miR-182-5p, and hsa-miR-26a-5p for the substantial presence of BCR-ABL1 in CML adults treated with TKIs at the molecular response. Additionally, we have mRNA NGS data from our CML patients treated with TKIs. The study objective is to demonstrate the miRNA-mRNA interactions and thus verify the three validated miRNAs as biomarkers for monitoring CML treated with TKIs. Methods: mRNA NGS data was analysed using bioinformatics analysis. Targets of the three validated miRNAs were predicted using molecular prediction tools and databases. Results: Differential expression gene analysis between cases and controls showed that inframe insertion was observed only in non-responsive patients. Interestingly, insertions included three similar genes that were PPIP5K2, POLE3, and NADK. Duplication was observed with Lysine on PPIP5K2 and POLE, while Glutamic acid on NADK. These genes were listed on the Proteomic and Genomic Data Commons (PDC/GDC), a repository for tumors and cancer. Moreover, prediction tools predicted these genes were among the targets of the validated miRNAs. Thus these verified the interactions between the three validated miRNAs and mRNA in CML patients treated with TKIs. Conclusion: These showed that the three miRNAs could be used as new biomarkers in monitoring CML to complement current detection methods. Further analysis of the interactions will

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Polygenic Risk Score Significantly Predicted Coronary Heart Disease Death among those with Type 2 Diabetes in The Malaysian Cohort

^{1,2}Khairul Anwar Zarkasi, ¹Nor Azian Abdul Murad, ^{1,3}Noraidatulakma Abdullah, ⁴Norfazilah Ahmad, and ¹Rahman Jamal*.

¹*UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia*

²*Biochemistry Unit, Faculty of Medicine and Defence Health, Universiti Pertahanan Nasional Malaysia.*

³*Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia*

⁴*Faculty of Health Sciences, Universiti Kebangsaan Malaysia.*

ABSTRACT

Polygenic risk scores (PRS) have been widely researched to measure the risk of coronary heart disease (CHD), however, they are mainly optimised for the Caucasian population. There is a lack of PRS specifically targeting the Malaysian population. This study was aimed at developing a PRS to predict CHD death risk among type 2 diabetes (T2D) participants in The Malaysian Cohort project (TMC). In Phase I, T2D TMC participants (224 “T2DM+CHD” as cases, 625 “T2DM” as controls) were genotyped using genomic microarray followed by genetic imputations using GenomeAsia reference panel to identify single nucleotide polymorphisms (SNP) associated with CHD death and to construct the PRS. In Phase II, T2D participants without known CHD during recruitment (n=806) and followed-up until their deaths, were analysed based on their PRS risk categories. The outcome of interest was CHD deaths in both phases. From the 56 CHD-related SNPs with suggestive genome-wide significance identified in Phase I ($P < 5.00 \times 10^{-5}$), PRS was constructed using 15 SNPs genotyped from the microarray. Compared to “T2DM”, the “T2DM+CHD” had higher PRS scores across ethnicities while the area under the receiver operating curve (AUC) showed the ability of PRS to classify CHD death. Using the cut-off point of 0.8493 (AUC=0.762, sensitivity=64.7%, specificity=73.1%), the odds of CHD death in the high-risk category were 6.96, 7.90, and 22.58 for Malay, Chinese, and Indian, respectively. In Phase II, survival analysis showed that mean survival years were shorter in high-risk than the low-risk category among Malays (6.60 vs 8.92 years), Chinese (6.20 vs 8.66 years), and Indians (7.21 vs 8.60 years). Meanwhile, among high-risk participants, the CHD death risk was 3.32, 3.36, and 12.72 times higher among Malays, Chinese, and Indians, respectively. The potential use of PRS to predict CHD death among T2D patients could assist in the long-term management and risk identification, subsequently enhancing the precision medicine practice in Malaysia.

Keywords: CHD death; coronary heart disease; type 2 diabetes mellitus; single nucleotide polymorphism; polygenic risk score; The Malaysian Cohort Project

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Revealing Uncharted Depths of the Dark Proteome: A Re-examination of Mass Spectrometry Data Uncovers New Insights into Colorectal Cancer Microproteins

¹Leong, Alyssa Zi-Xin; ²Nur Arifah Syahirah Rosli; ¹Lee, Pey Yee; Mohtar, ¹Mohamad Aimanuddin
*¹Low, Teck Yew

¹ UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Kuala Lumpur, 56000, Malaysia.

² Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, 43400, Malaysia.

* lowteckyew@ppukm.ukm.edu.my

ABSTRACT

Background: Cancer proteogenomic projects like the Clinical Proteomics Tumor Analysis Consortium (CPTAC) concentrate on identifying and quantifying those ~20,000 proteins sourced from annotated open reading frames (ORFs). However, these initiatives inadvertently overlook non-canonical proteins emerging from unconventional ORFs. Nonetheless, increasing experimental evidence has pointed to previously inactive regions that now yield steady, functional proteins—spanning alternate proteins (with over 30 amino acids) and their subset, microproteins (below or equal to 100 amino acids). Methods: We downloaded a dataset of TMT-10plex MS-based proteomics data CPTAC acquired from colorectal cancer (CRC) tissues (PDC000116). We then re-analyzed this dataset using FragPipe harboring MSFragger, Philosopher, IonQuant and TMT-Integrator. MS data were searched against an in-house protein FASTA comprising human SwissProt, OpenProt and in silico translated, verified RIBO-Seq sequences. FragPipe Analyst was used for subsequent statistical analysis. Results: In PDC000116, we identified 10,634 proteins from all 100 samples, whereby 9482 were canonical proteins and 501 were alternate proteins. Out these 501 non-canonical proteins, 205 were ≤ 100 amino acids. Divided by the stages of CRCs, we found 10, 13, 10 and 18 upregulated microproteins for Stage 1, 2, 3 and 4 CRC respectively, whereas 34, 31, 27, 21 downregulated ones. Particularly, IP_613259, IP_650627 and IP_573056 were found to be significantly upregulated in all stages of cancer compared to normal. Conversely, IP_671454, IP_690979 and IP_624921 was found to be significantly downregulated in all cancer stages. Also, members of the S100 protein family were found in different stages of CRC, such as S100P, S100A8, S100A12, S100B and S100A1. Conclusion: The analysis revealed differentially expressed novel microproteins and alternate proteins, across colon cancer stages, highlighting their potential significance in cancer development.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Demethylbelamcandaquinone B from *Marantodes pumilum* var. *alata* (Blume) Kuntze inhibits osteoclast differentiation in RAW264.7 cells

Haryati Ahmad Hairi¹, Jamia Azdina Jamal², Nor Ashila Aladdin², Khairana Husain², Noor Suhaili Mohd Sofi³, Norazlina Mohamed³, Isa Naina Mohamed³, Ahmad Nazrun Shuid^{4*}

¹ Department of Biochemistry, Faculty of Medicine, Manipal University College Malaysia, Jalan Batu Hampar, Bukit Baru, 75150 Melaka, Malaysia

² Faculty of Pharmacy, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

³ Department of Pharmacology, Faculty of Medicine, Preclinical Building, Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, Bandar Tun Razak, Cheras, 56000 Kuala Lumpur, Malaysia

⁴ Department of Pharmacology, Faculty of Medicine, Universiti Teknologi Mara, Jalan Hospital, 47000 Sungai Buloh, Selangor, Malaysia

ABSTRACT

Background: Bone remodeling is stably maintained by an intricate balance between bone-forming osteoblast and bone-resorbing osteoclast activities. However, excessive bone resorption by osteoclast leads to bone loss diseases such as osteoporosis, periodontitis, rheumatoid arthritis, and bone cancer. Osteoclast bone-resorbing cells are characterized by positive tartrate-resistant acid phosphatase (TRACP) staining, which are originally derived from hematopoietic precursor cells of the monocyte/macrophage lineage. The present study was aimed at determining the effects of *M. pumilum* var. *alata* crude aqueous extract and its active compounds on osteoclast activity and cytokine secretion in osteoclast precursor cells induced by RANKL potentially through the oestrogen receptor (ER) signaling pathway. Method: RAW264.7 macrophages were differentiated using RANKL into osteoclast-like cells. Then, they were treated with 10 µg/mL *Marantodes pumilum* var. *alata* crude aqueous extract, 5 µg/ mL dichloromethane fraction, and 0.6 µg/mL Dmcq B and 0.06 µg/ mL estradiol. TRACP staining, TRACP 5b colometric assay, and bone-resorbing pits were examined. The gene and protein expression of pro-inflammatory cytokines (TNF-α and IL-6) and protein expression of ERα and ERβ were measured. Results: *Marantodes pumilum* var. *alata* crude aqueous extract and Dmcq B inhibited RANKL-stimulated osteoclast differentiation as evidenced by size reduction of giant multinucleated osteoclast cells, decreased TRACP 5b activity as well as the subsiding of resorbed pit area compared with normal control. In addition, they reduced the gene and protein expressions of TNF-α and IL-6. *Marantodes pumilum* var. *alata*, Dmcq B, and estradiol treatments increased the protein expressions of ERα and ERβ in osteoclasts. Conclusion: *Marantodes pumilum* var. *alata* and its active compound, Dmcq B can inhibit osteoclast differentiation.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

**GPC3 alterations and interactions in lung adenocarcinomas:
a comprehensive data mining study of publicly available datasets**

Shing Cheng Tan

*UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Cheras, Kuala Lumpur,
Malaysia*

ABSTRACT

Background: Glypican-3, encoded by the GPC3 gene, is a membrane-bound heparan sulfate proteoglycan commonly implicated in human lung adenocarcinomas (LUADs). Nevertheless, the function, mutational profile, epigenetic regulation, co-expression profile, and clinicopathological significance of the GPC3 gene in LUAD progression are not well understood. In this study, we aimed to elucidate the above parameters by mining cancer microarray datasets from publicly available databases. **Methods:** Oncomine, GEPIA2, and UALCAN databases were used for analysis of the GPC3 expression and methylation. The correlation between GPC3 expression and patient survival was then evaluated using PrognoScan. In addition, the correlation between GPC3 expression and immunological infiltration was investigated using TIMER. Mutational and CNA profiles of the GPC3 gene were analyzed using the cBioPortal platform. miRNAs and TFs interacting with GPC3 were identified using NetworkAnalyst v3.0. **Results:** We observed significant downregulation of GPC3 in LUAD tissues compared to their normal counterparts, and this downregulation was associated with shorter overall survival (OS) and relapse-free survival (RFS). Nevertheless, no significant differences in the methylation pattern of GPC3 were observed between LUAD and normal tissues, although lower promoter methylation was observed in male patients. GPC3 expression was also found to correlate significantly with infiltration of B cells, CD8+, CD4+, macrophages, neutrophils, and dendritic cells in LUAD. In addition, a total of 11 missense mutations were identified in LUAD patients, and ~1.4–2.2% of LUAD patients had copy number amplifications in GPC3. Seventeen genes, mainly involved in dopamine receptor-mediated signaling pathways, were frequently co-expressed with GPC3. We also found 11 TFs and 7 miRNAs interacting with GPC3 and contributing to disease progression. Finally, we identified three potential inhibitors of GPC3 in human LUAD, namely heparitin, gemcitabine and arbutin. **Conclusions:** GPC3 may play an important role in the development of LUAD and could serve as a promising biomarker in LUAD.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor

Phytochemical Profiling and DPPH (2,2-diphenyl-1-phyrcrilhydrazyl) Radical Scavenging Activities of The Crude Extract Ethanol *Laportea decumana* Rob. Wedd

Lukman La Basy^{1,2}, Triana Hertiani^{3*}, Retno Murwanti⁴, Ema Damayanti⁵

¹Pharmaceutical Sciences Doctoral Study Program, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

²Pharmacy Department, Stikes Maluku Husada, Maluku 97566, Indonesia

^{3*}Department of Pharmaceutical Biology, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia

⁴Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia

⁵Research Center for Food Technology and Processing, National Research and Innovation Agency (BRIN), Gunungkidul, 55861, Indonesia

ABSTRACT

Background; *Laportea decumana* Roxb. Wadd is an endogenous plant from the Maluku Islands that has been used topically by the local community for a long time as an analgesic. This study aims to determine the analgesic effect of the ethanol extract of itching leaves tested on mice *in vivo* and to determine the inhibition of Cox-2 enzymes *in vitro*, and determine the phytochemical profiling. Methods; the research material was obtained from Seram Island, Maluku, and the dry powder was extracted using 96% ethanol. The extract was carried out by phytochemical profiling and DPPH (2,2-diphenyl-1-phyrcrilhydrazyl) radical scavenging activities. Results; Phytochemical screening reveals the TLC profile of the crude extract containing flavonoids, phenolics, terpenoids, and alkaloids and then DPPH Radical Scavenging Activity of Crude Extract with an IC₅₀ value of 146.28 μ g/ml has moderate antioxidant activity. Conclusion; recommended for future studies of its pharmacological activity in providing evidence for its ethnomedical use.

Keywords: *Laportea decumana*; itch leaves; radical scavenging; Maluku.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Unraveling AGR2 Regulation in Breast Cancer: Insights from RNA-Seq Analysis of CRISPR/Cas9 mediated targeting.

**^{1*}M. Aiman Mohtar, ¹Syazalina Zahari, ¹Nisa Syakila A Rahman, ¹Saiful Effendi Syafruddin,
¹Muhammad-Redha Abdullah-Zawawi, ¹Low Teck Yew, ²Ted Hupp**

¹ UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, 56000 Kuala Lumpur, Malaysia

² Institute of Genetics and Cancer, MRC Institute of Genetics & Molecular Medicine, University of Edinburgh, Scotland EH4 2XR, United Kingdom

ABSTRACT

Anterior Gradient Protein-2 (AGR2), a member of the protein disulphide isomerase family, plays a pivotal role within the endoplasmic reticulum (ER) by managing the folding and maturation of client proteins as they navigate the secretory pathway. AGR2 has gained prominence as a key contributor to tumorigenesis and metastasis, with its upregulation observed in various cancer types, further fueling malignant progression. Interestingly, despite its primary residence in the ER, this intriguing protein possesses the ability to extend its influence beyond cellular confines and assume an oncogenic role in the extracellular environment. Nonetheless, the intricate mechanisms governing the functions of AGR2 and its secretion have remained enigmatic. Here, we unveiled the dual presence of AGR2 protein expression, both within breast cancer cells' intracellular compartments and in the extracellular space. Using the metastatic breast cancer cell line 1833-BoM as our experimental model, we have demonstrated that AGR2 secretion lacks glycosylation and partly involves unconventional protein secretion, and this was evidently pronounced under conditions of tunicamycin-induced ER stress. By employing lentivirus-delivered CRISPR-Cas9 technology to target AGR2, we have witnessed a reduction in cancer cell fitness, leading to impaired proliferation, adhesive capacity, invasion capabilities. Through comprehensive RNA-seq analysis, we have identified 603 significant differentially expressed genes in two isogenic AGR2-targeted cells. Functional enrichment analysis has shed light on the involvement of these genes in critical cellular processes, such as receptor signaling, oxidative stress response, cell adhesion, and vesicle trafficking. Further exploration via protein-protein interaction network analysis has revealed hub genes, including ESR1, AR, LCK, CDH1, and CD55, which have the potential to modulate AGR2 intracellular signaling and secretion. Our findings provide substantial evidence supporting the role of AGR2 in driving breast cancer progression, thereby highlighting its potential as a therapeutic target for interventions aimed at disrupting AGR2-associated pathways in breast cancer.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Integrative pipeline to analyze immune cells in digital pathology images

¹Haslina Makmur, ¹Pan Jia Wern, and ¹Teo Soo-Hwang*.

¹*Cancer Research Malaysia, Malaysia.*

ABSTRACT

Conventional digital pathology image analysis involves basic tasks such as cell detection, classification and quantification. While these may seem straightforward, their application to whole slide images is not simple, requiring intricate AI-driven image processing techniques that can be difficult for researchers to access. This study presents an integrative semi-automated pipeline to quantify immune cells on digital pathology images and derive predictive spatial features. Immunohistochemical (IHC) staining for CD3, CD4 and CD8 immune markers was performed on samples of breast cancer patients. After digitization, 43 images with a large tumour area were selected for further analysis. Immune cell detections were carried out on two different bioimaging software, QuPath and Fiji. Immune cells were detected, categorized and quantified in two areas (invasive front and tumour center) within tumour region using QuPath. Cell detection based on color threshold and region of interest (ROI) was conducted using Fiji. The ROI were divided based on their predicted area sizes into two levels – individual cells and groups of cells. The filtered ROI were used as inputs for CytoMAP to calculate distances between immune markers and local density of immune markers at both levels. Combining both results, a total of 48 spatial features derived from IHC immune markers were used to train a machine learning model to identify spatial features that were predictive of gene expression-based immune scores. Based on the machine learning prediction, we determined that the best predictive spatial features were the number of CD4 hotspots in tumor center and the distance between CD8 and other markers. Our results also suggest that immune features within the tumour center are more predictive than within the invasive front. These results provide insights into the use of off-the-shelf imaging software tools to analyze digital pathology images to make useful predictions in lieu of more complicated image processing techniques.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

KCNN4 mutation presenting with non-immune hydrops haemolytic anaemia

¹Nur Mashitah Abdullah, ¹Loh C Khai, ¹Doris Lau, ^{1,2,3*}Adli Ali

¹Department of Pediatrics, Hospital Pakar Kanak-Kanak (HPKK), The National University of Malaysia, Kuala Lumpur, Malaysia

²Institute of IR4.0, The National University of Malaysia

³Infection and Immunology Health and Advanced Medicine Cluster, The National University of Malaysia, Cheras, Kuala Lumpur, 56000 Malaysia

ABSTRACT

Background: Dehydrated hereditary stomacytosis (DHSt) is a rare autosomal dominant hemolytic anaemia disorder. The two genes reported in DHSt are PIEZO 1 gene and KCNN4 gene. In DHSt, the gene mutations cause dysfunctional membrane protein with the imbalance of potassium and sodium influx, resulting in the loss of water that leads to red blood cell dehydration and hemolysis. To date, there is no case reported in Asia. Case description: We report a case of an 8-year-old boy with DHTs who presented with hemolytic anaemia and non-immune hydrops at birth with strong family history of hemoglobinopathy with G6PD deficient Viangchan phenotypes. Antenatally, he was diagnosed as fetal hydrops with polyhydramnios at 28-week gestational age. He was noted to have anaemia at birth with a haemoglobin level of 7.3 g/dL and reticulocytes of 14% and required two times packed cell transfusions during neonatal period. His initial laboratory investigations for hemolytic anemia showed negative results for TORCHES and parvovirus screening but a mild increase in osmotic fragility test. He had a strong family history of anaemia involving his maternal side and siblings. His mother and maternal aunt had a history of cholecystectomy at 24 and 16 years old respectively. His mother and elder brother and sister had anaemia and confirmed to be heterozygous G6PD Viangchan. His parents were non-consanguineous. He had hepatosplenomegaly with poor growth due to chronic hemolytic anemia and presented with intermittent episodes of crisis with headache and jaundice. His brain imaging revealed dural venous malformation in magnetic resonance imaging and partial venous obstruction in cerebral angiogram. A whole exome sequencing showed positive KCNN4 gene mutation. Conclusion: DHSt is a rare hereditary disorder with no specific treatment. The mainstay treatment of DHSt is supportive with pack cell transfusion when indicated. Splenectomy should be avoided in DHSt as it increases the risk of thrombosis.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Risk Factors of Long COVID in Klang Valley Malaysia: A Snapshot from the COVGEN study

Noraidatulakma Abdullah¹, Ying-Xian Goh¹, Azwa Shawani Kamalul Ariffin¹, Aisyatul Najihah Khuzaimi¹, Mohd Arman Kamaruddin¹, Joanna Ooi Su Min², Azmawati Mohammed Nawi³, Norfazilah Ahmad³, Mohd Rohaizat Hassan³, Anita Sulong⁴, Najma Kori⁵, Mohamed Faisal Abdul Hamid⁵, Sheela Bai A/P Pannir Selvam⁶ and Rahman Jamal^{1*} on behalf of the COVGEN Cohort Study Team

¹ UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia.

² Anesthesiology & Intensive Care Department, Hospital Canselor Tuanku Muhriz (HCTM), UKM, Malaysia.

³ Department of Community Medicine, Faculty of Medicine, UKM, Malaysia

⁴ Department of Microbiology and Immunology, Faculty of Medicine, UKM, Malaysia

⁵ Department of Medicine, Faculty of Medicine, UKM, Malaysia

⁶ Cheras Health Clinic, Cheras, Kuala Lumpur, Malaysia

*Correspondence: rahmanj@ppukm.ukm.edu.my; Tel.: +60-3-914500

ABSTRACT

Background: Long COVID is a condition where prolonged symptoms, after Covid-19 infection, that continue for at least three months after the initial infection and persisting for at least two months without any other explanation. A significant number of patients with COVID-19 experience this condition, however few studies have investigated the characteristic of this population, particularly in outpatient settings. Methods: We recruited 1,561 participants from the baseline infection who had previously contracted COVID-19 from the Klang Valley area, specifically at Hospital Canselor Tuanku Muhriz (HCTM), UKM Medical Molecular Biology Institute (UMBI), and Cheras district's COVID-19 Assessment Centre (CAC), from August 11, 2021, to September 30, 2022. We followed them up at 3, 6, and 12 months. Only participants who had at least one follow-up at any of these time points were included in this study. Among them, only 872 met the eligibility criteria for the final analysis. Self-reported questionnaires and blood samples were collected from participants both at baseline and during the follow-up sessions after obtaining their consent. Results: Out of 872 participants, 227 (26.0%) were still experienced prolonged symptoms. Most of the participants were aged 30-50 (59.5%), female (59.1%), and Malay (88.1%). More than 90% of the Long COVID participants completed two doses of vaccination and most of them received BNT162b2 vaccine. Having respiratory disease, high diastolic blood pressure, high haemoglobin A1c, and presenting with more than five COVID-19 symptoms during infection were significantly associated with long COVID. The most common symptoms that remained persistent were neurological and cognitive function, such as brain fog, memory problems, difficulty concentrating, decreased alertness, sleep issues, confusion, and speech difficulties regardless of whether it was 3, 6, or 12 months after their initial COVID-19 infection. While the most newly developed diseases due to COVID infection diagnosed by medical doctors were hypertension, cardiovascular disease, and diabetes. A phenomenon of diabetes regression was also observed during the 12 months follow-up among the Long COVID patient. Conclusion: Understanding the characteristics and effects of Long COVID will help facilitate the healthcare needs for the patients including monitoring and treatment.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

**Development of breast cancer spheroids as a model for a pre-clinical evaluation using
Vitamin C as Anti-cancer agent.**

Ali Saeed Ali Mussa^{1,2}, Khalid Mohamed Ali Hajissa², Noor Fatmawati Mokhtar³, Rohimah
Mohamud^{4*} and Rosline Hassan^{1*}

¹ Department of Haematology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian
16150, Kelantan, Malaysia; alimaha1341989@gmail.com; roslin@usm.my

² Department of Biology, Faculty of Education, Omdurman Islamic University, Omdurman PO. Box 382,
Sudan; khalid541983@yahoo.com

³ Institute for Research in Molecular Medicine (iNFORMM), Universiti Sains Malaysia, Health Campus,
Kubang Kerian 16150, Kelantan, Malaysia; fatmawati@usm.my

⁴ Department of Immunology, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian
16150, Kelantan, Malaysia; rohimahm@usm.my

ABSTRACT

Background: Breast cancer is a highly prevalent type of cancer that affects women globally, resulting in a significant number of fatalities. The treatment of breast cancer continues to pose significant challenges. Therefore, the development of effective therapeutic models is of utmost importance in order to improve treatment strategies and provide significant benefits to patients. Methods: MCF-7 and MDA-MB-231 were cultured as monolayers until they reached 70–80% confluency. A 96-well plate coated with agarose was utilized to generate a multi-cellular breast cell spheroid. The morphology of spheroids formation was examined and observed daily until day 3 under an inverted microscope and images were analyzed using ImageJ software (USA). The stock solution of vitamin C was diluted. The time and dose of Vitamin C were optimized. The CellTiter-Glo® 3D Cell Viability Assay (Promega, USA) and Annexin V-FITC/PI Apoptosis Detection Kit (BD, Franklin Lakes, NJ, USA) were employed to assess the cytotoxicity effect on the spheroids. Results: The results showed the formation of spheroids in MCF-7 and MDA-MB-231. Both exhibited unique morphological features that correspond to the behavior of these different types of breast cancer. Both apoptosis and cytotoxicity activities were higher in MCF-7 compared to MDA-MB-231 with vitamin C treatment. Conclusion: The spheroids models were developed and characterized. High-dose Vitamin C showed the ability to induce cytotoxicity. This highlights the usefulness of the breast cancer spheroid as a preclinical model for drug screening platforms.

Keywords:

Breast cancer; Spheroid; Vitamin C; Apoptosis; development

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Dipeptidyl peptidase 4 (DPP IV) inhibitory activity of leaf extract and sesquiterpene lactones isolated from *Smilax sonchifolius*

1,2Hady Anshory Tamhid, 1Triana Hertiani*, 1Yosi Bayu Murti, 1Retno Murwanti

1Faculty of Pharmacy, Universitas Gadjah Mada (UGM), Yogyakarta, 2Department of Pharmacy, Universitas Islam Indonesia (UII), Yogyakarta.

ABSTRACT

Background : The Sesquiterpene lactones, enhydrin and uvedalin, are the chemotype and subtype compounds of the yacon leaf. They are known to have anti-diabetic activity, but their mechanisms are not clear. One of the mechanisms of anti-diabetic agents is to inhibit the activity of the dipeptidyl peptidase 4 enzyme (DPP IV). The research aimed to determine the inhibition of the extract, enhydrin, and uvedalin isolated from the yacon leaf against the DPP IV enzyme activity. **Methods :** Yacon leaves were extracted using 70% ethanol by the maceration method. Enhydrin and uvedalin were obtained from previous research. The inhibitory activity of the DPP-4 enzyme was then determined by the fluorescence assay method using a multi-well plate reader. Sitagliptin was used as standard inhibitor of DPP IV enzyme. **Results :** The ethanol extract of yacon leaves inhibits the DPP IV enzyme with an IC₅₀ of 846.02 ppm. It contained 1.33 and 1.06% of the enhydrin and uvedalin compounds, respectively. The percentage inhibition of the DPP IV enzyme by the extract and uvedalin at a concentration of 250 ppm was 32.66 and 37.03%, respectively, while enhydrin showed no inhibitory effect. **Conclusion :** The inhibitory activity of the DPP IV enzyme in yacon leaf extract might be confirmed by the content of sesquiterpene lactones. In the study of these compounds, only uvedalin has inhibitory effects with low activity, so there may be other compounds that may be involved in its inhibitory activity. However, it may provide a new approach to the treatment of type 2 diabetes.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor

A KLF6-dependent PDGFB-mTOR signalling loop links lipid homeostasis and tumor growth in renal carcinoma

Saiful E. Syafruddin^{1, 2}, Paulo Rodrigues², Erika Vojtasova², Saroor A. Patel², Dora Bihary², and Sakari Vanharanta²

¹UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Jalan Yaacob Latiff, Bandar Tun Razak, Kuala Lumpur, 56000, Malaysia

²MRC Cancer Unit, University of Cambridge, Hutchison/MRC Research Centre, Box 197, Cambridge Biomedical Campus, Cambridge, CB2 0XZ, UK

ABSTRACT

Clear cell renal cell carcinoma (ccRCC) is characterized by frequent inactivation of the von Hippel-Lindau tumor suppressor (VHL) and consequent accumulation of hypoxia-inducible factor 2 alpha (HIF2 α), which drives renal tumorigenesis. Despite the genetic basis of ccRCC has been well-characterized, advanced ccRCC is associated with high mortality. Super enhancers have been reported to support the expression of critical transcriptional regulators in various biological contexts. To identify novel molecular mechanisms supporting ccRCC pathogenesis, we analysed H3K27ac ChIP-Seq data from ccRCC samples and looked for super enhancer-associated transcription factors. One of the strongest super enhancers was in the vicinity of KLF6, a zinc finger DNA binding transcription factor. CRISPR/Cas9 and CRISPRi-mediated perturbation of KLF6 in several VHL mutant ccRCC cell lines impaired cell growth in vitro and in vivo as well as reduced metastatic lung colonization. Transcriptomic analyses revealed that KLF6 depletion led to downregulation of several lipid homeostasis pathways downstream of SREBF1 and SREBF2, suggesting a role for KLF6 in regulating mTOR activity. In accordance, we found that KLF6 modulated mTORC1 activity in ccRCC by transcriptionally regulating the expression of PDGFB, an activator of the PIK3-AKT-mTORC1 signalling pathway. Targeting PDGFB in ccRCC led to impaired mTORC1 activity, and supplementing the KLF6-depleted cells with human recombinant PDGFB rescued mTORC1 activity. Finally, we further demonstrated that ccRCC cells-derived PDGFB is able stimulates mTORC1 activation in autocrine and paracrine manner. Collectively, our findings unravel the link between super enhancer-driven transcriptional networks and essential metabolic pathways may provide clues to the mechanisms that maintain the stability of cell identity-defining transcriptional programmes in cancer.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Advancements in Amber Suppression Technology: A Window into Molecular Interactions of Human Viruses.

Nur Firdaus Isa*, Surdensteeve Peter, Nur Amirah Mohd Rozali

Department of Biotechnology, Kulliyah of Science, International Islamic University Malaysia (IIUM), Jalan Sultan Ahmad Shah, Bandar Indera Mahkota, 25200 Kuantan, Pahang Malaysia.

ABSTRACT

Amber suppression technology, inspired by genetic code expansion (GCE), is a molecular tool that facilitates the precise incorporation of noncanonical amino acids into a site-specific target protein within cellular environments. The technique requires several components: a target gene, photoactivatable unnatural amino acids, and an amber suppressor tRNA (tRNACUA)/aminoacyl tRNA synthetase (aaRS) orthogonal pair. A primary challenge associated with this technology is that it involves the delicate task of optimizing a combination of key components to maximize successful gene code expansion. Here, we demonstrate that amber suppression technology unveils crucial residues within a Herpes Simplex Virus 1 (HSV-1) viral protein that photocrosslink with a host protein complex. This interaction implies that HSV-1 viral protein may be involved in inhibiting the transcription elongation of cellular genes, potentially serving as a pivotal factor in HSV-1 pathogenesis. Building upon the successful optimization of this molecular tool, we are currently extending its application to various other human viruses, including SARS-CoV-2, H5N1, and HIV-1. In conclusion, our research underscores recent advancements that emphasize the promising potential of amber suppression technology in a wide range of medical biotechnological applications.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Metabolomic Fingerprinting and Footprinting analyses on TNBC sh-TINCR cell lines

¹Mohd Ikhwan Ismail, ²Afreena Afiqah Azman, ³Mohd Izwan Yusof, ¹Fazleen Haslinda MH,
²Ezanee Azlina MH

¹Faculty of Pharmacy, Universiti Teknologi MARA (UiTM), Malaysia.

²UKM Medical Molecular Biology (UMBI), Universiti Kebangsaan Malaysia (UKM), Malaysia

³Faculty of Applied Science, Universiti Teknologi MARA (UiTM), Malaysia.

ABSTRACT

47.9% breast cancer cases in Malaysia were detected at late stage which 29% of the cases represented by the triple negative breast cancer (TNBC) subtype. The affected pathway of the long noncoding RNA (TINCR), which promotes proliferation and metastasis of cancer cells is poorly understood. In this study the TINCR-knocked-out TNBC cell lines were subjected to metabolomic analysis with the aim to identify key metabolites and discover metabolic pathways responsible for the poor prognosis of this cancer subtype. MDA-MB-231 (n=3) and HS578T (n=3) cells were knocked down via lentiviral delivery of shRNA vector. Cell lysates and culture medium were harvested and subjected to NMR 600 MHz analysis. NMR spectra were calibrated, baselined, bucketed and spectra bin file was exported to Metaboanalyst 5.0 for statistical analysis. Spermine and Nicotinic acid metabolites in MDA-MB-231 cell while Cystathionine and Pyridoxal metabolites in HS578T cell were found to be perturbed by the knocked-out process. This suggests that biosynthesis of cofactors metabolic pathway (KEGG:hsa01240) took place in the TNBC cell lines progression. Downstream analysis was done using protein-protein and gene-gene interactions database indicated that 2 proteins/genes to be the main factor in the process which are PNP and BHMT gene. Further analysis on these 2 genes could have a potential to shed some light to the discovery of prognostic biomarkers and effective target for therapeutic intervention for TNBC cancer patients.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Evaluation and Optimisation of Adipogenic Differentiation Protocol for the Human Adipocyte Tissue-derived Stem Cells.

¹Siti Aishah Sulaiman* and ¹Viceswarry Dorairaj

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia (UKM)

*Corresponding author: sitiaishahsulaiman@ukm.edu.my

ABSTRACT

Background: The in vitro adipocyte cell line model is often used to provide insight into fundamental aspects of adipocyte physiology. Although most studies used murine adipose cells, human adipose tissue-derived stem cells (ASC) isolated from various adipose depots have noteworthy translational significance and relevance for adipocyte mechanistic studies. However, the differentiation method using the ASC cell line is inconsistent across different studies. This study aimed to evaluate and determine the optimum media composition for mature adipocyte differentiation from the ASC cell line. **Methods:** Undifferentiated hTERT-immortalized human adipose-derived stem cell line (ASC52Telo, ATCC) was grown in four different adipogenic differentiation media for 18-21 days. The media composition was 1) M1: ATCC premade differentiation media, 2) M2: Published basic cocktail media without vitamins and hormones, 3) M3: Published complete cocktail media without fetal bovine serum (FBS), 4) M4: Modified M3 media supplemented with FBS and glutamax, and 5) stem cell media (undifferentiated, CON). Adipocyte differentiation was confirmed by the formation of lipid vesicles stained by Oil Red O. Expressions of adipocyte markers were measured via the Real-time quantitative PCR method. **Results:** M4 media have the highest differentiated mature adipocytes (92.7%) compared to CON (0%), M1 (45.0%), M2 (10.3%), and M3 (38.6%) (P-value <0.0001). Expressions of three adipocyte markers (FABP4, SREBP1C, and PPARG2) were increased following M4 media treatment compared to the CON (P-value <0.001). For M1 media treatment, only FABP4 marker expression was higher (P-value <0.05). The stem cell marker THY1 expression was reduced only following M4 media treatment compared to the CON (P<0.0001). These marker expressions did not change after the M2 and M3 media treatment. **Conclusion:** This study provides the optimum media composition for mature adipocyte differentiation from the human ASC cell line. Establishing optimum in vitro adipogenic differentiation protocol is vital for the future fundamental research on obesity.

Keywords: adipose tissue-derived stem cells (ASC), adipogenesis, differentiation, optimisation, obesity.

Acknowledgment: This study is sponsored by the Fundamental Research Grant Scheme (FRGS) from the Ministry of Higher Education (FRGS/1/2019/SKK08/UKM/03/8).

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

SLC6A14 Expression and Methionine Uptake Increased in Advanced Colorectal Cancer Cells Stage

^{1,2}Hazwani Mohd Yusof*, ²Sharaniza Ab-Rahim, ³A Rahman A Jamal and ²Musalmah Mazlan

¹*Department of Biochemistry, Faculty of Medicine, Manipal University College Malaysia, Jalan Batu Hampar, Bukit Baru, Melaka. 75150 Malaysia.*

²*Department of Biochemistry and Molecular Medicine, Faculty of Medicine, Universiti Teknologi MARA, Campus Sungai Buloh, 47000 Sungai Buloh, Selangor, Malaysia.*

³*UKM Medical Molecular Biology Institute, UKM Medical Centre, Jalan Yaacob Latif, Bandar Tun Razak, 56000 Cheras, Kuala Lumpur, Malaysia*

ABSTRACT

Backgrounds: Data from our previous metabolomics study reported that L-methionine was significantly increased in the more advanced stages of colorectal cancer cells (CRC). Upregulated amino acids transporter in cancer cells may affect the uptake of methionine into cells. In this study, we aim is to determine the association between the SLC6A14 expression and methionine uptake in CRC at different stages. **Methods:** Normal colon cell lines (CCD 841 CoN) and CRC cell lines; HT 29 (early stage) and HCT 116 (late stage) were cultured to determine the expression of SLC6A14 by western blot analysis. SLC6A14 inhibits with alpha-methyltryptophan (α -MT), 4.5 mM for 72 hours was performed. The viability of cells was also determined by MTS assay, and apoptosis assay. The L-methionine level was measured with and without inhibition of α -MT by L-methionine fluorescence assay. The data was analysed using GraphPad Prism Version 6. **Results:** The results show SLC6A14 expression level in CRC cells at early and late stages were significantly upregulated compared to normal. The inhibition of SLC6A14 with α -MT (4.5 mM) leads to significant decreased in cell viability and associated with increased apoptosis activity and decreased intracellular level of methionine. **Conclusion.** This study conclude that increased level of methionine observed in the previous metabolomics study is due to increase expression of SLC6A14 in CRC cells that is involved in the uptake of methionine into cells for their growth and proliferation.

Keywords: colorectal cancer, SLC16A4, methionine, stages.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Optimization of Annealing Temperature for Dengue Virus Serotyping Using Multiplex Polymerase Chain Reaction (PCR) Technique.

Nurul Syafiqah Nabilah Abdul Kadir¹, Siti Norlia Othman², Norzuriza Mohd Rais², Hui-min Neoh¹, Meng Ling Moi³, Najma Kori⁴, Petrick Periyasamy⁴ & Nor Azila Muhammad Azami¹.

¹UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia,

²Department of Medical Microbiology and Immunology, Hospital Canselor Tunku Muhriz, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

³School of International Health, The University of Tokyo, Tokyo, Japan

⁴Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

ABSTRACT

Malaysia is a dengue hyper-endemic country with all four serotypes of dengue virus (DENV) circulating simultaneously. All four serotypes of DENV have similar clinical manifestations, but the homology among serotypes is less than 80% at the amino acid level. Dengue serotyping using polymerase chain reactions (PCR) technique is widely used to determine the serotype of DENV due to fast turnover time. Multiplex PCR techniques were selected due to their feasibility in detecting all four serotypes of DENV in a single-tube reaction compared to the conventional PCR technique for DENV serotyping. In this study, we aimed to determine the optimum annealing temperature for multiplex PCR for all four primer sets to simultaneously detect all four serotypes of DENV. We used the universal forward primer (UFP DENV) and four different dengue virus-specific reverse primers (D1-TR1DENV1, D2-TR2DENV2, D3-TR3DENV3, D4-TR4DENV4), each with a final concentration of 200 μ M. Viral RNA of DENV-1, DENV-2, DENV-3, and DENV-4 cultured in BHK cells were extracted using SpinStar Nucleic Acid Extraction Kit 1.0. Then, cDNA was synthesized using Super Script III First-Strand Synthesis Super Mix and PCR was performed using EasyTaq PCR reagent. Gradient PCR was performed to determine the optimum annealing temperature using two sets of temperatures: 55°C to 60°C and 58°C to 60°C. The results show that the optimum annealing temperature for DENV-1, DENV-2 and DENV-3 was 59.2°C with an apparent band size of 482 bp for DENV-1, 119 bp for DENV-2 and 290 bp for DENV-3. However, we were not able to get any amplification for DENV-4. Thus, we aimed to design a new primer for DENV-4. In conclusion, the optimum annealing temperature to be used in the multiplex PCR was 59.2°C. Therefore, we decided to use this temperature for validation with clinical samples.

Keywords: dengue virus, serotyping, multiplex PCR, annealing temperature, optimization

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Current status of predictive serum biomarkers for polyps and colorectal cancer

Maryam Fatimah Abu Bakar¹, Azmawati Mohammed Nawi^{1*}, Chin Siok Fong², Suzana Makpol³

¹Department of Public Health Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia.

²UKM Medical Molecular Biology Institute (UMBI), UKM Medical Centre, 56000, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia.

³Department of Biochemistry, Faculty of Medicine, Universiti Kebangsaan Malaysia, 56000, Jalan Yaacob Latif, Bandar Tun Razak, Kuala Lumpur, Malaysia.

ABSTRACT

Early detection of colorectal cancer (CRC) is crucial to enhance the disease treatment and prognosis of patients. Colonoscopy remains a golden method for CRC detection which requires trained personnel with expensive tools. Other non-invasive methods available such as faecal occult blood test (FOBT), however, still have a problem with high false positive rates. Currently, serum metabolites biomarker has been discovered to be potentially used to discriminate patients with polyps and CRC. This systematic review focuses on identifying the most commonly detected predictive serum metabolites for polyps and colorectal cancer. A systematic search of the Web of Science, PubMed, and Cochrane Library databases was conducted using PRISMA guidelines. Nine studies investigating serum metabolite biomarkers of colorectal cancer and polyps using different analytical platforms and study populations were included. QUADOMICS tool was used to analyze the quality of included studies. All metabolites detected for polyps and CRC was enriched into pathways using MetaboAnalyst 5.0. Metabolites detected by three studies and more might be potentially used as a predictive biomarker for polyps and CRC. A review of these studies revealed several potential signature metabolites overlapped between studies including Tyrosine, Lysine, Cystine, Arabinose, and Lactate for CRC, and Lactate and Glutamate for polyps. The most affected pathways related to CRC were the urea cycle, glutathione metabolism, purine metabolism, glutamate metabolism, and ammonia recycling, while the urea cycle, glutamate metabolism, glutathione metabolism, arginine and proline metabolism, and carnitine synthesis were found to be affected in the polyps. However, the differences between altered pathways in polyps and CRC, other external factors, and their effects on the regulation level, sensitivity, and specificity of each identified metabolite still remained unclear. Further research is needed to validate this finding and would benefit from large cohort studies, standardized analysis instruments, and well-defined subject groups.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

High-throughput genotyping of Asian-specific genetic variants associated with colorectal cancer risk in the Malaysian population: A pilot study

Mohamad Ayub Khan Sharzehan¹, Shing Cheng Tan^{1*}

¹UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Cheras, Kuala Lumpur, Malaysia

*Corresponding author: sctan@ukm.edu.my

ABSTRACT

Background: Identification of genetic variants associated with colorectal cancer (CRC) risk has shown promise in preventing the disease. Although it is well-established that the influence of genetic variants on CRC risk varies by populations and environmental factors, the association of genetic variants with CRC risk in the Malaysian population has not been studied using high-throughput genotyping method. In this work, we aimed to conduct a pilot genome-wide association studies (GWAS) on Malays using a genotyping panel developed specifically for the Asian population, with the aim to identify genetic variants associated with CRC risk. Methods: A case-control study was performed on 95 CRC patients and 95 healthy controls. Genotyping of all samples was performed with the Illumina Infinium Asian Screening Array using DNA obtained from peripheral blood of the study subjects. Genetic data were first subjected to preliminary analysis using Plink and R software to identify the association between individual polymorphisms and CRC risk. Logistic regression was performed using SPSS to determine the significance of the polymorphisms, along with epidemiological data, in predicting CRC risk. Results: Our preliminary analysis identified 10 suggestive significant polymorphisms (rs76997593, rs79946793, rs10483660, rs77350431, rs79883457, rs10870101, rs12434275, rs7815453, rs6471236, and rs5747442) were statistically significant for CRC risk prediction in the genetic models. When epidemiological data (age, BMI, and marital status) were incorporated into the prediction model, only rs76997593 and rs10870101 maintained their significance with regard to CRC risk. However, the statistical significance diminished when TNM staging and histological grading were included in the prediction model. Conclusion: The results obtained suggests that rs76997593 and rs10870101 could serve as potential markers for predicting the risk of CRC in Malays, although further studies with a larger sample size are required to clarify this point.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Novel Ryr2 receptor gene mutation presenting with cardiac tumour and refractory arrhythmias: Case report and systemic review

¹Nur Mashitah Abdullah, ^{1,2,3*}Adli Ali

¹*Department of Pediatrics, Hospital Pakar Kanak-Kanak (HPKK), The National University of Malaysia, Kuala Lumpur, Malaysia*

²*Institute of IR4.0, The National University of Malaysia*

³*Infection and Immunology Health and Advanced Medicine Cluster, The National University of Malaysia, Cheras, Kuala Lumpur, 56000 Malaysia*

ABSTRACT

Background: Ryanodine Receptor 2 (RYR2) gene mutation causing catecholaminergic polymorphic ventricular tachycardia (CPVT) is one of the identified causes of sudden death in adult and children. Case description: We report a case of novel Ryanodine Receptor 2 (RYR2) gene mutation presented with cardiac arrest and recurrent syncopal attack with accidental finding of cardiac tumour. Methods: Case was reported based on retrospective review of information via electronic medical record, digital laboratory system, and digital radiological, image and reports in Hospital Pakar Kanak-Kanak (HPKK), Universiti Kebangsaan Malaysia. For systemic review, three large databases; PubMed, Ovid and Scopus were used to find articles with keywords “RYR2 gene mutations” and “CPVT” up to December 2022. Selected articles based on title and abstract included are: (i) case reports and articles on children 18 years old and below with (ii) treatment response on CPVT and (iii) written in English only. We excluded reports of asymptomatic patients with RYR2 gene mutations and articles written in other languages. Results: 14 studies were chosen and reviewed together with our reported patient. Most of the patients presented initially with syncopal attack and developed cardiac arrest later. Some of them presented with both syncopal attack and seizures precipitated by exercise or stress. We found that 43.8% of patients shared similar variants or coding effects in RYR2 gene mutation. Demographically, the mean age at presentation is 11 years old with 53% of reported cases were male. Conclusions: Refractory arrhythmias with cardiac arrest not responded to adrenaline should raise the suspicion towards RYR2 gene mutations. Recognition of this condition is important as it affects the outcome of resuscitation. Untimely diagnosis of RYR2 gene mutations with appropriate use of pharmacological agent during resuscitation is important to ensure a better outcome.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Whole exome sequencing (WES) in Mullerian Agenesis and Anorectal Anomaly: A Case Study

Siti Aishah Sulaiman^{1*}, Chow Yock Ping¹, Muhammad-Redha Abdullah Zawawi¹, Zam Zureena Mohd Rani¹, Siti Nurmi Nasir¹, Salwati Shuib², Dayang Anita A. Aziz³, Hana Azhari³, Sharifah Azween Syed Omar⁴, Zarina Abdul Latiff⁴, Rahman Jamal¹, Nor Azian Abdul Murad^{1*}

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia (UKM), Jalan Yaacob Latif, 56000, Cheras, Kuala Lumpur.

²Department of Pathology, ³Department of Surgery, ⁴Department of Pediatric, Faculty of Medicine, Universiti Kebangsaan Malaysia (UKM), Jalan Yaacob Latif, 56000, Cheras, Kuala Lumpur.

*Corresponding author: nor_azian@ppukm.ukm.edu.my & sitiaishahsulaiman@ukm.edu.my

ABSTRACT

Background: Mullerian agenesis or Mayer-Rokitansky-Kuster-Hauser Syndrome (MRKH) Type II is a congenital malformation of the Mullerian ducts that results in the absence of a uterus in women. The aetiology of this syndrome is unknown, and it is considered a sporadic genetic disease. MRKH, together with anorectal anomaly (VACTERL), is extremely rare. Several cases have yet to be reported so far without any information on genetic analysis. This study determined the mutation profile of a girl diagnosed with Mullerian Agenesis and Anorectal Anomaly with Rectovaginal Fistula. Methods: Whole-exome sequencing (WES) was performed on a 5-year-old Malaysian girl diagnosed with Mullerian Agenesis and Anorectal Anomaly with Rectovaginal Fistula and her normal parents. Data were analyzed using Torrent Suite v.5.0.4 and annotated using ANNOVAR. Single nucleotide polymorphisms (SNPs) with an allele frequency > 0.01 were excluded, and the remaining variants were filtered based on de novo mutations and autosomal recessive and autosomal recessive inheritance traits. Related genes were then analyzed to determine the enrichment of function and pathway sites (g:Profiler) and protein-protein interactions (HIPPIE v. 2.3, STRING v.11.5, and GeneMANIA). Results: Mullerian Agenesis disease in this patient may be caused by 28 de-novo mutations. Two mutations in the LHX5 (p.P358Q), inherited from the father and CFTR (p.R1158X), inherited from the mother, are likely to be associated with the MRKH syndrome in this patient. Conclusion: A total of two inherited mutations and 28 de-novo mutations in 30 genes have been identified, and interaction with each other is likely causing MRKH syndrome in this patient.

Keywords: Rare disease-Mullerian Agenesis and Anorectal Anomaly, Whole Exome Sequencing

Acknowledgment: This study is sponsored by the UKM Genomic Unit HCTM-UMBI.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Integrin Beta 1/CD29: a preliminary investigation into a possible GBM EV surface marker

Johannes Low Jun Wei and Nadiah Abu*

UKM Medical Molecular Biology Institute, Kuala Lumpur.

ABSTRACT

Background: Glioblastoma multiforme (GBM), is both malignant and hard to detect yet extracellular vesicles, facilitates disease detection via liquid biopsy. In this study, we elucidated exosomal surface epitopes of GBM clinical and cell lines' EVs and offered insights into a potential GBM EV surface marker. **Methodology:** GBM derived EVs were collected from clinical (5 GBM patients vs 5 healthy controls) and cell line (A172, LN18, U251MG, KNS42) settings via the precipitation and size exclusion chromatography combination strategy. After BCA quantitation, isolated EVs characterized for their particle size (Zetasizer), morphology (transmission electron micrography) and surface exosomal epitopes (MACSPlex Exosome Kit, human). 6 µg EVs were incubated with capture beads before staining with APC-conjugated antibody cocktails, followed with flow cytometry detection. Positive surface epitopes were run through RNA and proteomics dataset to determine their RNA and protein level in the cell. Literature search was performed to determine literature relevant to epitope expression. **Result:** GBM derived EVs displayed typical round morphology within 30 -200 nm exosomal range. MACSPlex analysis revealed varying exosomal surface epitopes expression for both clinical and cell lines' EVs: CD63 being the most expressed tetraspanins for clinical EVs while CD81 is most prominent among GBM cell lines' EVs. Interestingly, GBM markers were among the exosomal surface epitopes list (CD29, CD44 and CD146). Clinical EVs displayed higher expression of CD146 compared to healthy controls, while cell line EVs has higher expression of CD29 and CD44. Downstream analyses on positive epitopes revealed ITGB1 as a cell surface protein on U-251MG, LN-18 and ex vivo GBM sample (Cell Surface Protein Atlas) while Human Protein Atlas, ProCan-DepMapSanger and CCLE mass spectrometry data revealed ITGB1 is more expressed in RNA and protein level across tested GBM cell lines **Conclusion:** ITGB1 as a GBM invasion driver is also expressed on GBM EV surface.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Radioactive Iodine Refractory Papillary Thyroid Cancer: ABC Transporter Genes Under the Microarray Lens

¹Nur Fadhina Mohamad Pakarulrazy, ²Nadzlee Harith Paisol, ²Rohaizak Muhammad, ²Shahrin Niza Abdullah Suhaimi, ³Nurismah Md Isa, ¹Nadiah Abu, ¹Nurul Syakima Ab Mutalib, and ²Nani Latar*.

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Kuala Lumpur;

²Department of Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur; ³Department of Pathology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur.

ABSTRACT

The mortality rates in cases with papillary thyroid cancer (PTC) are notably elevated when characterised by lymph node positivity (LN+) and resistance to radioactive iodine (RAI-R) ablation. ATP-binding cassette (ABC) transporters, a class of transmembrane proteins, have been closely associated with drug resistance. However, the relationship and roles of these transporters in LN+ and RAI-R PTC have not been determined. To address this knowledge gap, our study aimed to identify differentially expressed ABC transporter genes in RAI-R PTC using microarray analysis. A total of five thyroid tumour tissues without lymph node metastasis (LN-) and with RAI avidity (RAI-A), five thyroid tumour tissues with LN+ and RAI-R, and four adjacent normal thyroid tissues were collected from PTC patients operated at Hospital Canselor Tuanku Muhriz UKM (HCTM). Histopathological evaluations were performed on the collected tissues before the extraction of total RNA. Subsequently, we conducted mRNA profiling using the Agilent SurePrint G3 Human Gene Expression v3 8x60K Microarray. Finally, AltAnalyze and NetworkAnalyst software were utilised to conduct probeset expression analysis. The defined criteria for the identification of differentially expressed ABC transporter genes are p-value < 0.05 and fold change (FC) ≥ 2 or ≤ -2 . In the analysis of 47 ABC transporter genes, it was observed that three genes, namely ABCB4, ABCB7, and ABCF2, exhibited differential expression exclusively in RAI-R PTC. The expression of ABCB4 (FC = 2.41) and ABCB7 (FC = 2.03) genes were significantly upregulated in RAI-R PTC compared to normal thyroid tissue. In contrast, the expression of the ABCF2 gene was significantly downregulated in RAI-R PTC relative to both normal thyroid tissue (FC = -2.79) and RAI-A PTC (FC = -2.44). In conclusion, our findings suggest that the ABCB4, ABCB7, and ABCF2 genes may play pivotal roles in RAI-R PTC. However, comprehensive investigations are warranted to elucidate their involvement in thyroid cell iodine uptake.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

CovGen Study: Dissecting COVID-19 Symptom Severity and Risk Factors

¹Nor Adzimah Johdi, ¹Noraidatulakma Abdullah, ¹Nor Azila Muhammad Azami, ¹Ying-Xian Goh, ¹Azwa Shawani Kamalul Ariffin, ¹Aisyatul Najihah Khuzaimi, ¹Mohd Arman Kamaruddin, ²Joanna Ooi Su Min, ³Azmawati Mohammed Nawi, ³Norfazilah Ahmad, ³Mohd Rohaizat Hassan, ⁴Anita Sulong, ⁵Najma Kori, ⁵Mohamed Faisal Abdul Hamid, ⁶Sheela Bai A/P Pannir Selvam and ¹Rahman Jamal* on behalf of the COVGEN Cohort Study Team

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia.

²Anesthesiology & Intensive Care Department, Hospital Canselor Tuanku Muhriz (HCTM), UKM, Malaysia.

³Department of Community Medicine, Faculty of Medicine, UKM, Malaysia

⁴Department of Microbiology and Immunology, Faculty of Medicine, UKM, Malaysia

⁵Department of Medicine, Faculty of Medicine, UKM, Malaysia

⁶Cheras Health Clinic, Cheras, Kuala Lumpur, Malaysia

*Correspondence: rahmanj@ppukm.ukm.edu.my; Tel.: +603-91459000

ABSTRACT

The COVID-19 pandemic, stemming from coronavirus SARS-CoV-2, has emerged as a profound global health crisis. This study investigates the association between risk factors and the severity of COVID-19 symptoms. Conducted between August 2021 and October 2022, this study encompassed laboratory-confirmed COVID-19 patients admitted to the Hospital Canselor Tuanku Muhriz, Universiti Kebangsaan Malaysia in Cheras, Kuala Lumpur. We classified stage I to III as mild, while stage IV and V as severe, according to ISARIC - WHO Case Report Form. Among the 554 participants, 413 (74.5%) were admitted to general wards, with 223 (97.4%) encountering mild symptoms and 190 (58.5%) facing severe symptoms. Notably, 141 (25.5%) required ICU admission, with 6 (2.6%) displaying mild symptoms and 135 (41.5%) experiencing severe symptoms. Hospitalization demonstrated a strong association with symptom severity ($p < 0.001$), with an adjusted odds ratio of 15.36 for ICU admission ($p = 0.009$). Age also held significance ($p < 0.001$), as those over 55 exhibited an adjusted odds ratio of 2.85 ($p = 0.009$) for severe symptoms. Ethnicity emerged as a significant factor ($p < 0.001$), with Malay, Indian, and other groups exhibiting milder symptoms compared to the Chinese patients. Comorbidities such as hypertension and obesity showed a notable link to severe symptoms ($p = 0.005$, $p = 0.034$ respectively). Vaccination demonstrated a mitigating effect on symptom severity while the vaccine brand correlated with symptom severity. The Delta variant showed greater severity compared to the Omicron variant (adjusted OR = 4.05, $p = 0.005$). Symptoms like systemic, cardiovascular, musculoskeletal, and certain neurological symptoms indicated higher risk ($p < 0.05$). Abnormal laboratory parameters—ALT, total protein, albumin, LDH, lymphocyte count, neutrophil count, CRP, and D-dimer—linked to severe symptoms ($p < 0.05$). Elevated HbA1c levels were tied to greater severity ($p = 0.030$, OR=1.10), while lower calcium levels indicated higher risk ($p < 0.001$, OR=0.02). This study underscores the multifaceted factors influencing COVID-19 symptom severity, offering insights for patient management and tailored interventions based on risk profiles.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

High Glucose Level Reduces Cisplatin Chemosensitivity in Cervical Cancer Cells

¹Siti Aisyah Md Sanib, ¹Rehanna Mansor*, ²Wing Hin Lee, ³Nur Najmi Mohamad Anuar

¹*Cluster of Integrative Physiology and Molecular Medicine, Universiti Kuala Lumpur Royal College of Medicine Perak, Ipoh, Perak, Malaysia.*

²*Faculty of Pharmacy and Health Science, Universiti Kuala Lumpur Royal College of Medicine Perak, 30450 Ipoh, Perak, Malaysia*

³*Programme of Biomedical Science, Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur 50300, Malaysia*

*Corresponding author email: rehannam@unikel.edu.my

ABSTRACT

Cervical cancer is the fourth most common cancer in women worldwide. Recent evidence suggests that metabolic syndrome and diabetes may influence cancer prognosis. Studies have shown that patients with cervical cancer and pre-existing diabetes have a worse prognosis than those without. Currently, chemotherapy is the main treatment for inoperable or advanced postoperative cervical cancer. The effectiveness of cisplatin to treat advanced or recurring cervical cancer may, however, be severely hampered by the development of cisplatin resistance. The aim of this study is to investigate the effect of high glucose condition on the response of cervical cancer cells to chemotherapy. Using Hela cells, we investigated the effect of different glucose conditions: 5-, 9- and 25-mM glucose on cells' response to chemotherapy drug cisplatin at concentration of 10 μ M and 60 μ M. Trypan Blue dye-exclusion assay was used to determine the cell death. We found a significant reduction of cisplatin-induced cell death on Hela cells grown in 9 mM and 25 mM glucose conditions compared to Hela cells grown in normal glucose conditions (5mM). This finding suggested that high glucose condition induces resistance to chemotherapy drugs in cervical cancer cells. These findings may indicate the importance of personalized treatment and management of cervical cancer patients with pre-existing diabetes.

Keyword: Hyperglycaemia, Cervical Cancer, cisplatin, chemoresistance

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Chemical Characterization, Antioxidant Potential and in vitro Anticancer properties of Rice (*Oryza sativa* var Bajong) from Sarawak

¹Lam Xiu Qian, ¹Isabel Fong Lim*, ²Khong Heng Yen*, ¹Tay Siow Phing

¹Department of Paraclinical Sciences, Faculty of Medicine and Health Sciences, Universiti Malaysia Sarawak,
Kota Samarahan 94300, Sarawak, Malaysia

²School of Chemistry and Environmental Studies, Faculty of Applied Sciences, Universiti Teknologi MARA,
Sarawak branch, Kota Samarahan 94300, Sarawak, Malaysia

* Correspondence: flisabel@unimas.my; khonghy@uitm.edu.my

ABSTRACT

Background: Rice (*Oryza sativa* L.) is an important staple food consumed by almost half of the world's population. Many consumers prefer polished white rice and neglect the nutritional benefits contained in pigmented rice. While many studies had reported on the nutritional and medicinal properties of pigmented rice bran layer, there is a knowledge gap regarding rice germ and endosperm layer. This is the first time the bran, germ and endosperm layer of darkly pigmented *Oryza sativa* var Bajong (Bajong) of Sarawak were assessed for its chemical, antioxidant and anti-cancer properties against human colorectal cancer cell line HCT-116 respectively. Methods: Bajong was extracted over 24-, 48- and 72-hour durations in alcohol. The extracts were characterised chemically for their total phenolic content, total flavonoid content and antioxidant potential respectively. The anticancer effects of Bajong extracts on HCT-116 cells were determined by 3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide (MTT) cell viability assay and Annexin-V apoptosis assay respectively. Results: The results showed Bajong was rich in total phenolic and flavonoid contents, exhibited strong free-radical scavenging activity, efficacious in promoting antiproliferation and triggering apoptosis in HCT-116 cells in dose-dependent and time-dependent manner respectively. The 72-hour Bajong extract exhibited the strongest growth inhibition in HCT-116 cells than those of 48-hour extract. Additionally, it significantly triggered high population of HCT-116 cells (51.2%) to undergo apoptosis at 72 hours. Conclusion: *Oryza sativa* var Bajong is rich in polyphenols and flavonoids which are pertinent and evident by its strong antioxidant activities. It showed promising efficacy in significantly reducing HCT-116 cell viability and triggering apoptosis in dose- and time-dependent manners. These findings showed Bajong possessed high potential as nutraceutical crop and promising chemopreventive agent against colorectal cancer.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

LRG1 Induces Lysosomal Autophagy via Dysregulation of the mTOR Pathway

^aLim SM, ^aMohamad Hanif EA, ^aChin S-F*

a UKM Medical Molecular Biology Institute, National University of Malaysia, Kuala Lumpur, Malaysia

ABSTRACT

Leucine-rich-alpha-2-glycoprotein 1 (LRG1) has emerged as a key player in various malignancies, including colorectal cancer (CRC). LRG1 exhibits a stage-dependent expression in CRC patients, and it correlates with tumour size. Several functions of LRG1 have been revealed in CRC such as proliferation, migration and invasion. Recently, LRG1 has been reported to regulate autophagy during cerebral ischemia/reperfusion injury and acute myocardial infarction. However, the autophagic role of LRG1 in CRC and its signalling pathway remains elusive. Hence, genetic modulation of LRG1 was conducted in CRC cells, HT-29 and COLO320DM to study the function of LRG1 in autophagy. HT-29 was overexpressed with LRG1 while LRG1 depletion was done in COLO320DM. The autophagy activity of the LRG1-modulated CRC cells was measured by evaluating the gene and protein expression of autophagy markers, LC3B and p62, through qPCR, western blot, immunofluorescence and flow cytometry. The activity of mTOR pathway was determined by gene expression of mTOR pathway components (Akt, mTOR and Raptor) and phosphorylation of ribosomal protein S6 (RPS6). Overexpression of LRG1 increased the transcript level of p62, but not LC3B, in HT-29 cells. The protein expression of autophagy markers, LC3B and p62, was concomitantly increased with LRG1 overexpression, suggesting the induction of impaired autophagy. Furthermore, LRG1 enhanced the gene expression of mTOR pathway components and phosphorylation of RPS6, indicating activation of the mTOR pathway. In contrast, LRG1 depletion upregulated the gene expression of LC3B but not p62. Surprisingly, silencing LRG1 promoted the protein expression of both LC3B and p62, similar to the overexpression model. However, LRG1 knockdown suppressed the gene expression of mTOR pathway components and phosphorylation of RPS6, implying the inhibition of mTOR pathway. Taken together, genetic modulation of LRG1 gene in CRC cells impacted on the mTOR pathway, although both overexpression and knockdown resulted in activation of impaired autophagy.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

**Abstract for 9th Regional Conference on Molecular Medicine (RCMM)
in Conjunction with 5th National Conference for Cancer Research (NCCR)
14-15th October 2023, Bangi Resort Hotel, Selangor**

Chemosensitization of TINCR: potential key regulator to chemoresistance in triple negative breast cancers (TNBCs).

¹Ezanee Azlina Mohamad Hanif*, ¹Afreena Afiqah Azman, ¹Chin Siok Fong, ²Nor Fadilah Rajab, ³Rohaizak Muhammad, ³Nani Harlina Md Latar, ⁴Reena Rahayu Md Zin.

¹UKM Medical Molecular Biology Institute (UMBI), Universiti Kebangsaan Malaysia, Kuala Lumpur, ²Centre for Healthy Aging & Wellness, Universiti Kebangsaan Malaysia, Kuala Lumpur, ³Faculty of Medicine (Surgery Department), UKM Medical Centre, Universiti Kebangsaan Malaysia, Kuala Lumpur, ⁴Faculty of Medicine (Pathology Department), Universiti Kebangsaan Malaysia, Kuala Lumpur.

ABSTRACT

Terminal Differentiation-Induced Non-coding RNA (TINCR) proven to be involved in molecular/cellular processes in triple negative breast cancer (TNBC). High TNBC recurrence is common. Therefore, this study aimed to assess TINCR level of expression across breast cancer cell panel and evaluate its prognostic relevance in TNBC. Additionally, to assess chemosensitivity to chemotherapy drugs and migration effects in TINCR-transduced TNBC cells. QPCR was carried out on a panel of 8 breast cancer cell lines (184A1, MCF10A - Normal-like; T47D, MCF7 – Luminal; MDA-MB-468, HCC1143 – Basal A TNBC; Hs578T, MDA-MB-231 – Basal B TNBC). Kaplan-Meier survival curve of TINCR was obtained from KM plotter database to indicate the prognostic values of high/low TINCR status. Two TNBC cells (MDA-MB-231 and Hs578T) were stably transduced via lentiviral transduction of shNT and shTINCR plasmids deliveries. The stably-transduced cells were then subjected to drug response curve (DRC) of chemo-drug treatments (5-fluorouracil (5-FU), epirubicin, cyclophosphamide, FEC; 5-FU, Epirubicin, Cyclophosphamide combination and Taxotere) followed by MTT assay for EC50 determination. TINCR expression was found highly expressed in Basal B TNBC cells compared to other cells. Survival analysis indicated shorter survival in patients with elevated TINCR expressions (HR = 2.69; p=0.0044). shTINCR cells displayed higher chemosensitivity to 5-FU (EC50= 11.95; p= 0.0001), Cyclophosphamide (EC50= 11.58; p= 0.03), and Taxotere (EC50= 7.13; p<0.0001) in MDA-MB-231 cells compared to MDA-MB-231-shNT. Hs578T-shTINCR chemosensitized cyclophosphamide (EC50= 9.55; p=0.0007) Taxotere (EC50=0.0537) and FEC (EC50=0.00128; p<0.0001). Reduced migration rate (n=1) were seen in 5-FU, Cyclophosphamide, epirubicin and taxotere in MDA-MB-231-shTINCR compared to shNT. These preliminary data indicated that TINCR has a potential role in chemosensitizing chemo-drugs which may also play part in the migration of cells in tumour setup. By observing DRC individually and combination (FEC), drug efficacy is greater in combination drug plus TINCR inhibition, thus suggesting minimal toxicity in treatment management which however require further assessments.

IN CONJUNCTION WITH THE
5TH NATIONAL CONFERENCE FOR
CANCER RESEARCH

*THE CONFERENCE THAT BRINGS TOGETHER PROFESSIONALS AND EXPERTS
FROM VARIOUS FIELDS TO DISCUSS THE LATEST DEVELOPMENTS AND TRENDS IN
PRECISION MEDICINE IN HEALTHCARE*

UKM MEDICAL MOLECULAR BIOLOGY INSTITUTE (UMBI)

*JALAN YA'ACOB LATIFF,
BANDAR TUN RAZAK, 56000 CHERAS,
WILAYAH PERSEKUTUAN KUALA LUMPUR*



For Further Enquiry:

✉ rcmm.umbi@gmail.com

☎ 03-9145 9070 / 03-9145 9223

<https://www.ukm.my/umbi/rcmm-2023/>